

# Functional analysis reveals Miocene fossil pig's foraging behavior

July 28 2014




Fig.1 Rostral and masticatory muscles of *Chleuastochoerus*. (a) lateral view of skull; (b) ventral view of skull; (c) lateral view of mandible. Scale bars equal 5 cm. Credit: HOU Sukuan

Foraging behavior and adaptations for feeding in the context of habitat

condition have been studied widely in living suids (pigs) but rarely in their fossil relatives. A study published in *Science China: Earth Sciences* (5) compared the skull and mandible of a Late Miocene fossil pig from the Linxia Basin, Gansu Province, China, *Chleuastochoerus*, with those of extant pigs and peccaries, and revealed the foraging behavior of *Chleuastochoerus*, adding fresh evidence that the area of the Linxia Basin was open but still relatively humid in the middle-late Late Miocene.

*Chleuastochoerus* is a small Asian endemic Late Miocene fossil pig, and its geographic range may be limited to northern China and the southwestern part of Asiatic Russia. In recent years, abundant specimens of *Chleuastochoerus* have been found in the Linxia Basin, Gansu Province, China. These new finds have included skulls and mandibles of juveniles and adult males and females, and even a nearly complete skeleton. These new specimens provide an opportunity for more detailed studies of the genus.

The [skull](#) of *Chleuastochoerus* is less elongated than in extant suids, and its dorsal profile has only a slight posterodorsal inclination unlike in extant suids and tayassuids (peccaries). "The presence of a rostral bone in the skull and relatively strong rostral muscles indicates that *Chleuastochoerus* possessed considerable digging ability, though the relatively narrow occipital surface and the shortness of the skull would have restricted the range of motion through which the skull could be swung", said lead author Dr. HOU Sukuan, Institute of Vertebrate Paleontology and Paleoanthropology of the Chinese Academy of Sciences (IVPP), "The distinctive pre-zygomatic plate and over-canine arch-niche may have played a role in protecting the skull and canines during digging".

The massive masticatory muscles of *Chleuastochoerus*, reconstructed based on well-developed parietal and zygomatic crests, would have

enabled the mouth to close quickly and contributed to moving the mandible both longitudinally and laterally. The configuration of the cranio-mandibular joint and its relatively high position show clearly that Chleuastochoerus was capable of both crushing and grinding movements during mastication.




Fig.2 Left mandibles of Hylochoerus (a), Phacochoerus (b), Sus (c), and Chleuastochoerus (d) in occlusal view. Scale bars equal 3 cm. Credit: HOU Sukuan

The dentition of Chleuastochoerus is generally Sus-like, but simpler and more primitive. The incisors of Chleuastochoerus are clearly primitive.

The comparatively simple structure of the crown surfaces of the cheek teeth suggests that the food of *Chleuastochoerus* may have been softer than that of extant suids. The diet of *Chleuastochoerus* may have been intermediate in composition between those of true forest suids (*Sus*, *Potamochoerus*) and open habitat suids (*Hylochoerus*, *Phacochoerus*) today, perhaps most closely resembling that of the savannah-adapted *Potamochoerus*, and the habitat of *Chleuastochoerus* should be relatively humid forest edge or nearby areas of open steppe.

This work was supported by National Natural Science Foundation of China, Chinese Academy of Sciences, Ministry of Science and Technology of China, and Key Deployment Project of Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences.

**More information:** "Foraging behavior of *Chleuastochoerus* (Suidae, Artiodactyla) A case study of skull and mandible morpho-functional analysis." HOU SuKuan, et al. *SCIENCE CHINA Earth Sciences* 2014 57 (5): 988-998 ISSN: 1674-7313 CN: 11-5843/P

Provided by Chinese Academy of Sciences

Citation: Functional analysis reveals Miocene fossil pig's foraging behavior (2014, July 28) retrieved 22 September 2024 from

<https://phys.org/news/2014-07-functional-analysis-reveals-miocene-fossil.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.