


A sponge-like molecular cage for purification of fullerenes

December 15 2014, by Daniel Maspoch, Inhar Imaz



A work in *Nature Communications* presents a supramolecular nanocage which encapsulates fullerenes of different sizes and allows the extraction of pure C₆₀ and C₇₀ through a washing-based strategy. The work was coordinated from the Universitat de Girona. Contributors include Ramón y Cajal Researcher Dr. Inhar Imaz and ICREA Research Prof. Daniel Maspoch from the ICN2 Supramolecular NanoChemistry & Materials Group.

A fullerene is a molecule made of carbon that can be found in many shapes, such as spheres or tubes. They have a wide range of applications, highlighting their extensive use as electroactive materials in solar cells

and with continuously appearing new uses in medicine. Any application, however, is limited in origin by tedious solid-liquid extractions (usually in toluene) and time-expensive chromatographic separations. Even the extraction of small amounts of purified fullerenes for research purposes becomes a complex process.

A work in *Nature Communications* recently presented a supramolecular nanocage synthesized by metal-directed self-assembly, which encapsulates fullerenes of different sizes. The study was led from the Universitat de Girona and included the participation of the UAB Department of Chemistry, the ALBA Synchrotron and the ICN2 Supramolecular NanoChemistry & Materials Group. RyC Researcher Dr. Inhar Imaz and ICREA Research Prof. Daniel MasPOCH are among the authors.

The article provides direct experimental evidence for the 1:1 encapsulation of C_{60} , C_{70} , C_{76} , C_{78} and C_{84} , fullerene molecules consisting of 60, 70, 76, 78 and 84 carbon atoms respectively. Solid state structures for the caged fullerenes with C_{60} and C_{70} have been obtained using X-ray synchrotron radiation. With a washing-based strategy it is possible to exclusively extract pure C_{60} from a solid sample of cage charged with a mixture of fullerenes.

This tetragonal prismatic supramolecular cage with a high affinity for the inclusion of fullerenes, and a facile ability to release them by solvent washing of the solid inclusion compound, is an attractive methodology to selectively extract C_{60} and, with a lower efficiency, C_{70} from fullerene mixtures. Although the method cannot be used to produce big amounts of purified fullerenes, it provides an experimental platform to design tuned cages for selective extraction of higher [fullerenes](#). The solid-phase fullerene encapsulation and liberation represent a twist in host-guest chemistry for molecular nanocage structures.

More information: Cristina García-Simón, Marc Garcia-Borràs, Laura Gómez, Teodor Parella, Sílvia Osuna, Jordi Juanhuix, Inhar Imaz, Daniel Maspoch, Miquel Costas & Xavi Ribas. Sponge-like molecular cage for purification of fullerenes. *Nature Communications*, 2014. 5, 5557. [DOI: 10.1038/ncomms6557](https://doi.org/10.1038/ncomms6557)

Provided by Catalan Institute of Nanoscience and Nanotechnology

Citation: A sponge-like molecular cage for purification of fullerenes (2014, December 15) retrieved 6 September 2024 from <https://phys.org/news/2014-12-sponge-like-molecular-cage-purification-fullerenes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.