

Butterflies deceive ants using chemical strategies

April 8 2015



Photo of a 4th instar larvae of *Arhopala zylda* and *Crematogaster* ant workers on a *Macaranga beccariana* tree. Credit: Yoko Inui, Lambir Hills National Park, Malaysia

Oakblue butterflies may use a variety of chemical strategies to deceive ants and avoid their attack, according to a study published April 8, 2015 in the open-access journal *PLOS ONE* by Yoko Inui from Osaka Kyoiku University, Japan and colleagues.

Certain plants have a mutualistic association with ants and are generally well-protected from herbivore attacks by the ant. However, larvae of Oakblue butterflies survive and develop on specific plant, *Macaranga*, or ant-plant species (the host), without being attacked by the ants living on the plant (known as plant-ant species). The authors of this study hypothesized that the butterfly larvae may chemically mimic or camouflage themselves and be accepted by the plant-ants on their host plant, and therefore would not be accepted by other plant-ant species living on [plants](#) that are not their normal host species.

To understand this process, the scientists collected butterfly larvae for three *Arhopala* species in Malaysia and then analyzed the plant-ant species behavioral and chemical responses to the experimental introduction of butterfly larvae, as well as larval dummies treated with cuticular hydrocarbon extract, to the leaves of *Macaranga* species. Cuticular hydrocarbons are chemicals that play an important role in insect communication.

The experiments revealed that although the responses of the plant-ants to the [butterfly larvae](#) varied considerably depending on the butterfly

species, attacks by plant-ant [species](#) on the normal plant host were infrequent. Chemically speaking, however, *A. dajagaka* matched well the host plant-ants, *A. amphimuta* did not match, and unexpectedly, *A. zylda* lacked hydrocarbons. Behaviorally, the larvae and dummies coated with cuticular chemicals of *A. dajagaka* were well attended by host and non-host plant-ants, *A. amphimuta* were often attacked by non-[host plant](#)-ants, and *A. zylda* were ignored by all [ants](#). The study authors suggest that variations exist in the chemical strategies used by the gossamer-winged butterflies that allow them to avoid ant attack and be accepted by the plant-ant colonies.

More information: Inui Y, Shimizu-kaya U, Okubo T, Yamsaki E, Itioka T (2015) Various Chemical Strategies to Deceive Ants in Three Arhopala Species (Lepidoptera: Lycaenidae) Exploiting Macaranga Myrmecophytes. *PLOS ONE* 10(4): e0120652. [DOI: 10.1371/journal.pone.0120652](#)

Provided by Public Library of Science

Citation: Butterflies deceive ants using chemical strategies (2015, April 8) retrieved 6 July 2024 from <https://phys.org/news/2015-04-butterflies-ants-chemical-strategies.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.