

West Antarctic Glaciers Are Increasingly Thinning

September 25 2004

Glaciers in West Antarctica are shrinking at a rate substantially higher than observed in the 1990s. They are losing 60 percent more ice into the Amundsen Sea than they accumulate from inland snowfall. The study was conducted by a science team from [NASA](#), U.S. universities and from the Centro de Estudios Científicos in Chile. It is based on satellite data and comprehensive measurements made in 2002 by a science team aboard a Chilean P-3 aircraft equipped with NASA sensors. Science Express published the findings today.

The ice loss from the measured glaciers corresponds to an annual sea-level rise of .008 inches (.2 millimeters) or more than 10 percent of the total global increase of about .07 inches (1.8 millimeters) per year.

About Antarctic Ice

Two ice sheets cover Antarctica, the larger East Antarctic Ice Sheet and the smaller West Antarctic Ice Sheet. The East Antarctic Ice Sheet is land-based, meaning that most of its base lies above sea level. The West Antarctic Ice Sheet is a marine-based ice sheet. The East Antarctic Ice Sheet is fairly stable and responds to environmental changes slowly. In contrast, the West Antarctic Ice Sheet may be capable of changing very rapidly.

For a balanced glacial system, the amount of glacier ice melting or flowing into the sea roughly equals the ice formed from snow accumulations further inland. The scientists report the Amundsen Sea glaciers are not in balance.

Bob Thomas, a science team member with EG&G Services at the NASA Goddard Space Flight Center's Wallops Flight Facility, Wallops Island, Va., commented that as the glaciers flow to the ocean, they become afloat to form ice shelves. "The ice shelves act like a cork and slow down the flow of the glacier," Thomas said.

"Ice shelves in the Amundsen Sea appear to be thinning, offering less resistance to their tributary glaciers. Our measurements show an increase in glacier thinning rates that affects not only the mouth of the glacier, but also 60 miles (100 kilometers) to 190 miles (300 kilometers) inland," Thomas said.

The scientists noted the earth underneath the ice is further below sea level than had been assumed, so the ice is thicker than once thought. This increases the amount of ice each glacier can discharge into the ocean as its speed increases. It makes it easier for the thinning glacier to float free from its bed, and thus further 'loosen the cork'," Thomas said.

Thomas pointed out the observed increases in velocities and thinning rates apply to only a short time period, so it is too early to tell if the accelerated thinning is part of a natural cycle or is a sign of a longer-term change. "Continued observation is important," he said.

"The rates of glacier change remain relatively small at present," said Eric Rignot, a study participant from NASA's Jet Propulsion Laboratory, Pasadena, Calif. "But the potential exists for these glaciers to increase global sea level by more than one meter. The time scale over which this will take place depends on how much faster the glaciers can flow, which we do not know at present," he said.

Thomas said in the last 10 years the ability to accurately measure glaciers worldwide has greatly improved. Measurements from aircraft and satellites like NASA's Ice, Cloud and Land Elevation Satellite

(ICESat), launched in 2003, have greatly improved accuracy.

Citation: West Antarctic Glaciers Are Increasingly Thinning (2004, September 25) retrieved 26 April 2024 from <https://phys.org/news/2004-09-west-antarctic-glaciers-increasingly-thinning.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.