

New twist on spin hall effect

February 8 2005

An international team of physicists that includes a Texas A&M University professor has announced discovery of a new spintronic effect in semiconductor chips, the intrinsic spin Hall effect, which puts a new twist on future technology and the possibility for novel circuits with low energy consumption.

The team is formed by physicists Dr. Jürg Wunderlich and Dr. Bernd Kaestner from the Hitachi Cambridge Laboratory, U.K.; Prof. Tomáš Jungwirth from the Institute of Physics of the Academy of Sciences of the Czech Republic and the University of Nottingham, U.K.; and Prof. Jairo Sinova from Texas A&M.

In a normal Hall effect, a voltage is created perpendicular to an electric current as it flows through a conductor in a magnetic field. The magnetic field deflects the moving charges to the sides of the conductor, resulting in an observable Hall voltage.

The spin Hall effect was first predicted in 1971. Here the moving electrons, which carry with them a tiny magnet called the "spin," collide with impurities and these collisions generate opposing magnetizations at the conductor's edges.

Despite its intriguing ramifications, the theory disappeared into virtual obscurity until 1999, when it was rediscovered and further elaborated. Four years later, two independent teams, one including Sinova and Jungwirth, proposed a novel mechanism called the intrinsic spin Hall effect in which similar magnetization occurs without the need for

collisions.

The prediction touched off a theoretical firestorm, resulting in more than 50 articles arguing for and against the possibility. As the heated debate raged on, Wunderlich and Kaestner developed a new type of device to measure magnetization at each side of a high-mobility, ultra-thin conducting layer embedded within a semiconductor chip using built-in light-emitting diodes.

Armed with this novel tool, Wunderlich and Kaestner teamed with Jungwirth and Sinova to observe the spin Hall effect. Their findings will be featured the February issue of Physics Today along with an independent and parallel observation of the effect in conventional bulk semiconductors.

Team members say the more than 10 times larger signal detected in the Hitachi device can be attributed to the special layered design of the semiconductor chip that yielded operation in a regime close to the intrinsic spin Hall effect.

"They are both beautiful experiments, and one of the most remarkable aspects is that they seem to be exploring opposite regimes," Sinova adds.

The possibility of generating magnetization without circulating currents has great implications in many areas, most notably the design of information processing and storage devices.

"Obviously we are only at the beginning of this journey of discovery," Sinova explains. "As you gather more facts, the truth tends to reveal itself. That's the fun of science. We're seeking to know, and we're learning in the process."

Citation: New twist on spin hall effect (2005, February 8) retrieved 25 April 2024 from <https://phys.org/news/2005-02-hall-effect.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.