

Large scale ozone losses

April 27 2005

Scientists from the EU SCOUT-O3 Integrated Project, which is co-ordinated by the University of Cambridge's Chemistry Department, have been studying the links between stratospheric ozone and climate change in the Arctic since May 2004. This recent finding was announced during a meeting of the European Geophysical Union in Vienna on Monday, 25 April.

Overall temperatures in the ozone layer were the lowest for 50 years and were consistently low for over three months. From late November to late February, large areas of polar stratospheric clouds (PSCs) - clouds in the ozone layer- were present over the Arctic region at altitudes between 14 and 26 km.

This is the largest in the 50 year record, and especially in the last 20 years, the period when the ozone-depleting compounds have been high.

The chemical balance in the stratosphere is changed significantly by the presence of these clouds, altering the breakdown products from manmade CFCs (chlorofluorocarbons) so that rapid chemical ozone destruction can occur in the presence of sunlight. The cold conditions affected the distribution of nitrogen oxides, allowing ozone loss to continue longer than usual.

The European scientists reported the first signs of ozone loss in January 2005. As sunlight returned to northern latitudes the rate of ozone depletion increased and rapid destruction of ozone occurred throughout February and March. In the altitude range where the ozone layer usually reaches its maximum concentration, more than half of the ozone was

lost. "Overall about 30% of the ozone layer was destroyed," said Dr Markus Rex from the Alfred Wegener Institute in Potsdam, Germany. "This largely prevented the normal seasonal increase of the thickness of the ozone layer during winter and led to a thinner ozone layer in Arctic spring compared to warmer years."

The overall degree of ozone loss this year was of similar magnitude to the record loss that was observed in the Arctic during winter 1999/2000. During late March the Arctic air masses drifted over central Europe and contributed to individual days of significantly increased UV-B radiation and sunburn risk in parts of Europe. The affected region reached as far south as northern Italy.

Emissions of ozone depleting substances are now largely banned worldwide by the Montreal protocol. As a first success of this milestone of international cooperation in environmental policies the atmospheric concentrations of CFCs started to decrease. But the atmospheric lifetime of these compounds is extremely long and the concentrations will remain at dangerously high levels for another half century.

Over the next few decades the fate of the Arctic ozone layer will mainly depend on the evolution of atmospheric temperatures at the altitude of the ozone layer. Over the past forty years the conditions there have become significantly colder.

"The cooling was particularly pronounced for the cold Arctic winters. Unfortunately these are the winters that result in large ozone losses. In 2005 the average extent of conditions cold enough for the existence of polar stratospheric clouds was four times larger than it has ever been in the sixties or early seventies of the past century," said Dr Rex.

This continuous cooling trend is qualitatively consistent with what would be expected as a result of increasing concentrations of greenhouse gases

in the atmosphere. However the coupling processes between climate change and temperatures in the polar ozone layer are complicated by feedback processes that are currently not sufficiently understood to make reliable predictions for the future.

"Our aim is to improve the predictions of future ozone and other stratospheric changes as well as the associated UV and climate impact," said Dr Neil Harris from the University of Cambridge, one of the coordinators of the project.

"Within SCOUT-O3 we have followed the meteorological conditions in the Arctic closely and a suite of atmospheric observations and model calculations was triggered on a very short notice. The Arctic ozonesonde station network started a campaign of coordinated measurements to monitor the chemical ozone destruction. ESA carried out additional measurements of the chemical composition of air in the Arctic ozone layer with the ENVISAT research satellite. The high flying research aircraft Geophysica made a deployment deep into Arctic air masses resulting in additional in-situ observations of key species."

Preliminary results from all these studies are being presented at the European Geophysical Union meeting in Vienna this week.

Source: Cambridge University

Citation: Large scale ozone losses (2005, April 27) retrieved 18 April 2024 from <https://phys.org/news/2005-04-large-scale-ozone-losses.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.