

Nanoscale Fiber Optics

July 14 2005

Lipid membranes as molds for doped cadmium chloride nanowires with light-conducting properties

One-dimensional nanoscopic structures such as nanowires are important building blocks for future miniature opto-electronic components. Swiss researchers have now developed a new method for the production of [nanowires](#); they use lipid membranes as "molds" and obtain high yields of cadmium chloride nanowires that behave as light conductors.

“Syntheses that use molecular molds have advantages”, explains Horst Vogel, “they are simple, work under mild conditions, and deliver unique, precisely defined nanostructures”. Vogel and his team selected phospholipid membranes as molds. Phospholipids consist of a water-friendly head group and water-repellent tail groups (hydrocarbon chains). In aqueous surroundings, they line up tail to tail into double-layered membranes. If these are dried carefully, stacks of membranes are formed, in which the head groups that point toward each other are separated by nanometer-wide water films. Some types of head groups are able to selectively bind certain positively charged ions. This is the basis of the Swiss researchers’ technique.

They took phospholipids with a preference for cadmium ions and produced membrane stacks with high cadmium concentrations. They then treated these with hydrochloric acid fumes. The hydrogen atoms from the hydrochloric acid forced the cadmium ions out of the binding sites and drove them into the water layer between the membrane layers. Here they combined with the negatively charged chloride ions from the

acid to form tiny cadmium chloride crystals, which then continued to grow into one-dimensional wires. Why only one dimension?

Beyond a certain crystal size, the lipid head groups, given a positive charge by the hydrogen atoms, begin to interact with the surfaces of the crystal which are rich in negatively charged chloride ions. The crystal thus cannot grow any further at those surfaces. Because of the special crystal structure, one of the crystal surfaces is protected from these interactions by water molecules, which are also a part of the crystal lattice. Here the crystal continues to grow. In this way, wires of up to 170 μm long and only about 100 nm in diameter are formed, which consist of a single continuous crystal.

If the hydrochloric acid vapor is augmented with hydrogen sulfide, the result is cadmium chloride nanowires that contain fluorescing cadmium sulfide nanocrystals. “If we irradiate one of these nanowires in the middle, we don’t only observe fluorescence at that one location but also at both ends”, says Vogel. “The nanowire conducts the light, just like an optical fiber.”

Source: Angewandte Chemie

Citation: Nanoscale Fiber Optics (2005, July 14) retrieved 10 April 2024 from <https://phys.org/news/2005-07-nanoscale-fiber-optics.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.