

A sharp boundary in Earth's layers

July 30 2005

A team led by a Brown University student shows a sharp boundary exists between the Earth's hard outermost shell and its more pliable layer beneath.

Geophysicists from Brown University in Providence, R.I., and the Massachusetts Institute of Technology in Boston say their findings suggest strong evidence temperature alone can't account for differences between the regions, which allow plate tectonics to occur.

Earth's cool, rigid upper layer, the lithosphere, rides on top of its warmer, more pliable neighbor, the asthenosphere, as a series of massive plates. Plates continuously shift and break, triggering earthquakes, sparking volcanic eruptions, sculpting mountains and carving trenches under the sea.

Catherine Rychert, a graduate student in Brown's Department of Geological Sciences, found a sharp dividing line between the lithosphere and the asthenosphere, according to data culled from seismic sensors sprinkled across the northeastern United States and southeastern Canada.

The findings are published in Nature.

Copyright 2005 by United Press International

Citation: A sharp boundary in Earth's layers (2005, July 30) retrieved 9 April 2024 from



https://phys.org/news/2005-07-sharp-boundary-earth-layers.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.