

Rainbands Offer Better Forecasts of Hurricane Intensity

August 8 2005



Scientists will soon begin one of the largest research projects ever undertaken to better understand dramatic, rapid changes in hurricane intensity. These changes have baffled forecasters for decades.

Atmospheric scientists from the University of Miami Rosenstiel School of Marine & Atmospheric Science (RSMAS), the University of Washington and the National Center for Atmospheric Research (NCAR) in Boulder, Colo., will participate in the new project, called the Hurricane Rainband and Intensity Change Experiment, or RAINEX.

RAINEX will study how the outer rain bands and inner eye of a hurricane interact to influence a storm's intensity.

"While great progress has been made in forecasting hurricane tracks, we need to improve forecasting of hurricane intensity," said Steve Nelson,

director of the National Science Foundation's (NSF) physical and dynamic meteorology program, which funds RAINEX. "Many factors affect the intensity of hurricanes," Nelson said. RAINEX scientists will investigate one of those factors: the interactions between hurricane rainbands and the eyewall. "From RAINEX, we will better understand the impact of rainbands on a hurricane's maximum winds," he said.

While researchers have studied the eye and outer rainbands of hurricanes extensively, "few, if any, experiments have ever examined these two components together and how their interaction might affect a storm's strength," said Shuyi Chen, a meteorologist and physical oceanographer at RSMAS and a RAINEX principal investigator. "The outer bands of a hurricane often have strong winds and lots of rain, and that can actually affect the overall intensity of a hurricane," she said.

RAINEX will study this interaction using data recorded from hurricane research flights. Starting on Aug. 15 and continuing through the remainder of this year's Atlantic hurricane season, two NOAA P3 aircraft, along with a U.S. Navy P3 aircraft, all equipped with Doppler radar, will fly simultaneously into hurricanes well before they threaten landfall.

The University of Washington and NCAR, will conduct research using airborne Doppler radar analysis. RSMAS will construct a state-of-the-art hurricane model using the data collected from the research flights.

"These flights can be turbulent, especially when we're penetrating a hurricane's rainbands," said NCAR scientist Wen-Chau Lee. "I think that's the wild card, the challenge of the experiment: to capture internal rainband structure and its interactions with the eye wall in those conditions."

"We hope to find an explanation for why a hurricane changes in

intensity, from the relationship between the inner and outer parts of the storm," said Robert Houze, an atmospheric scientist at the University of Washington and a RAINEX co-principal investigator. "These storms can jump up in intensity, or drop a full category in a day, a big challenge."

Flying in the hurricane's outer bands and into the eye wall, scientists aboard the aircraft will use sophisticated Doppler radar and GPS dropsondes to record wind speed and direction, temperature, humidity, atmospheric pressure and other critical data.

Much of what scientists currently know about the interaction between the outer rainbands and the eye wall of a hurricane comes from the state-of-the-art numerical models developed for hurricane research and prediction, which can provide very detailed information but may not be completely accurate.

Researchers need solid data to validate these models, they say. "We need to know whether or not our models are accurate, and the data we collect from RAINEX will give us the information we need," Chen said.

Once the data are collected, the researchers will all analyze and share this information with hurricane operational centers and national environmental prediction centers throughout the country, and the world.

Source: The National Science Foundation

Citation: Rainbands Offer Better Forecasts of Hurricane Intensity (2005, August 8) retrieved 18 April 2024 from <https://phys.org/news/2005-08-rainbands-hurricane-intensity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.