

Seafloor Creatures Destroyed By Ice Action During Ice Ages

October 19 2005

The ice ages made massive changes to the Earth's landscape. But what was happening below the ice in the oceans? Research by marine scientists reveals that it was a time of mass destruction as whole communities of animals were wiped out by ice sheets scouring the sea floor.

In the past it has been thought that these ecosystems somehow dodged extinction by recolonising from nearby habitats that escaped obliteration. But researchers at the National Oceanography Centre, Southampton (NOC) and the British Antarctic Survey (BAS) reveal a bleaker scenario.

Dr Sven Thatje, an ecologist at NOC has been working with geoscientists, Dr Claus-Dieter Hillenbrand and Dr Rob Larter at BAS examining one of the harshest environments on Earth - the Antarctic seafloor.

Writing in the October issue of Trends in Ecology & Evolution the scientists provide new evidence that suggests that seafloor organisms were either erased by the advance of ice sheets across the Antarctic continental shelf or starved to death as links in the food chain were broken by the permanent ice cover.

There would have been no refuge for shallower living animals further down the continental slope, as huge sediment slides would have buried them. Typically these ecosystems would have been made up of sponges, urchins, sea fan corals, and starfish.

Dr Thatje said: 'We show that during ice ages seafloor organisms emigrated to the deep sea - below the effects of the sediment slides and ice. From there, organisms may have invaded open marine shelters of the Antarctic shelf, which were not affected by the advance of ice masses. Or these animals may have recolonised the Antarctic shelf from the deep-sea during the warm period following each ice age.'

'Either way it is an impressive feat against the odds as the extreme cold means that these animals respond much more slowly to the destruction of their habitat than elsewhere in the oceans. They have lower metabolic rates that lower their growth and reproductive rates. Elsewhere in the oceans, a brisingid starfish would reproduce annually and live for ten years. In the Antarctic these starfish can reach 100-years-old but reproduce only once every ten years. This means that full community recuperation takes up to hundreds of years.'

Dr Claus-Dieter Hillenbrand explained: 'Until now it was commonly thought that the destructive action of the ice sheets was not significant enough to eradicate all the fauna and that desolate patches were recolonised from surrounding areas. But our research confirms that the destruction was wholesale with very little surviving. Even today calving icebergs ploughing across the seafloor destroy everything in their path. Imagine the impact of ice sheets during the ice ages that covered a much wider area in a time of lower sea levels.'

The team's research will lead to a radical rethink of the evolutionary history of Antarctica as the work challenges all the accepted theories. The scientists argue that shallow water animals were retreating to the deep ocean and then returning to recolonise Antarctica's shelf seas. Clues to how these two very different communities could have achieved this may lie in the animals' DNA.

Sven Thatje continued: 'Our work means that the text books will need to

be rewritten. Our next task is to reconstruct what happened in Antarctica during these periods of climate change and study the genetic and biological links between deep sea and shallow water communities.'

Copyright 2005 by Space Daily, Distributed United Press International

Citation: Seafloor Creatures Destroyed By Ice Action During Ice Ages (2005, October 19)
retrieved 18 April 2024 from
<https://phys.org/news/2005-10-seafloor-creatures-ice-action-ages.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.