

# Allo, allo? A star is ringing

December 21 2005

---



Astronomers have used ESO's Very Large Telescope in Chile and the Anglo-Australian Telescope in eastern Australia as a 'stellar stethoscope' to listen to the internal rumblings of a nearby star. The data collected with the VLT have a precision better than 1.5 cm/s, or less than 0.06 km per hour!

*Image: Alpha Centauri and the Southern Cross.*

By observing the star with two telescopes at the same time, the astronomers have made the most precise and detailed measurements to date of pulsations in a star similar to our Sun. They measured the rate at which the star's surface is pulsing in and out, giving clues to the density, temperature, chemical composition and rotation of its inner layers -

information that could not be obtained in any other way.

The astronomers from Denmark, Australia, and the USA [1] used Kueyen, one of the four 8.2-m Unit Telescopes of ESO's Very Large Telescope (VLT) at Cerro Paranal in Chile, and the 3.9-m Anglo-Australian Telescope (AAT) in New South Wales (Australia), to study the star Alpha Centauri B, one of our closest neighbours in space, about 4.3 light-years away.

Alpha Centauri is the brighter of the two 'Pointers' to the Southern Cross. Alpha Centauri itself is a triple system and Alpha Centauri B is an orange star, a little cooler and a little less massive than the Sun.

Churning gas in the star's outer layers creates low-frequency sound waves that bounce around the inside of the star, causing it to ring like a bell. This makes the star's surface pulsate in and out by very tiny amounts - only a dozen metres or so every four minutes [2].

Astronomers can detect these changes by measuring the small, associated wavelength shifts.

The researchers sampled the light from Alpha Centauri B for seven nights in a row, making more than 5 000 observations in all. At the VLT, 3379 spectra were obtained with typical exposure times of 4 seconds and a median cadence of one exposure every 32 seconds! At the AAT 1642 spectra were collected, with typical exposures of 10 s, taken every 90 s.

"From this unique dataset, we were able to determine as many as 37 different patterns (or modes) of oscillation", says Hans Kjeldsen, from University of Aarhus (Denmark) and lead author of the paper describing the results [3].

The astronomers also measured the mode lifetimes (how long the oscillations last), the frequencies of the modes, and their amplitudes

(how far the surface of the star moves in and out). Such measurements are a huge technical challenge. Indeed, the star's surface moves slowly, at the tortoise-like speed of 9 cm a second, or about 300 metre an hour. The astronomers borrowed their high-precision measurement technique from the planet-hunters, who also look for slight Doppler shifts in starlight.

"So much of what we think we know about the universe rests on the ages and properties of stars," said Tim Bedding, from the University of Sydney and co-author of the study. "But there is still a great deal we don't know about them."

By using two telescopes at different sites the astronomers were able to observe the Alpha Centauri B as continuously as possible.

"That's a huge advantage, because gaps in the data introduce ambiguity," said Bedding. "The success of the observations also depended on the very stable spectrographs attached to the two telescopes -- UVES at the VLT and UCLES at the AAT -- which analysed the star's light."

Source: ESO

Citation: Allo, allo? A star is ringing (2005, December 21) retrieved 9 April 2024 from <https://phys.org/news/2005-12-allo-star.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---