

Hubble 'weighs' Dog Star's companion

December 13 2005



For astronomers, it's always been a source of frustration that the nearest white-dwarf star is buried in the glow of the brightest star in the nighttime sky. This burned-out stellar remnant is a faint companion of the brilliant blue-white Dog Star, Sirius, located in the winter constellation Canis Major.

Image: Based on the Hubble measurements made with the Space Telescope Imaging Spectrograph, an international team found that Sirius B has a mass that is 98 percent that of our own Sun. Credit: ESA and NASA

Now, an international team of astronomers has used the keen eye of the



NASA/ESA Hubble Space Telescope to isolate the light from the white dwarf, called Sirius B. The new results allow them to measure precisely the white dwarf's mass based on how its intense gravitational field alters the wavelengths of light emitted by the star.



This picture is an artist´s impression showing how the binary star system of Sirius A and its diminutive blue companion, Sirius B, might appear to an interstellar visitor. Credit: NASA, ESA and G. Bacon (STScI)

"Studying Sirius B has challenged astronomers for more than 140 years," said Martin Barstow of the University of Leicester, U.K., who is the leader of the observing team. "Only with Hubble have we at last been able to obtain the observations we need, uncontaminated by the light from Sirius, in order to measure its change in wavelengths."

"Accurately determining the masses of white dwarfs is fundamentally important to understanding stellar evolution. Our Sun will eventually become a white dwarf. White dwarfs are also the source of Type Ia supernova explosions that are used to measure cosmological distances



and the expansion rate of the universe. Measurements based on Type Ia supernovae are fundamental to understanding 'dark energy,' a dominant repulsive force stretching the universe apart. Also, the method used to determine the white dwarf's mass relies on one of the key predictions of Einstein's theory of General Relativity; that light loses energy when it attempts to escape the gravity of a compact star."

Sirius B has a diameter of 12,000 kilometres, less than the size of Earth, but is much denser. Its powerful gravitational field is 350,000 times greater than Earth's, meaning that a 68 kilogram person would weigh 25 million kilograms standing on its surface. Light from the surface of the hot white dwarf has to climb out of this gravitational field and is stretched to longer, redder wavelengths of light in the process. This effect, predicted by Einstein's theory of General Relativity in 1916, is called gravitational redshift, and is most easily seen in dense, massive, and hence compact objects whose intense gravitational fields warp space near their surfaces.

Based on the Hubble measurements of the redshift, made with the Space Telescope Imaging Spectrograph, the team found that Sirius B has a mass that is 98 percent that of our own Sun. Sirius itself has a mass of two times that of the Sun and a diameter of 2.4 million kilometres.

White dwarfs are the leftover remnants of stars similar to our Sun. They have exhausted their nuclear fuel sources and have collapsed down to a very small size. Despite being the brightest white dwarf known, Sirius B is about 10,000 times fainter than Sirius itself, making it difficult to study with telescopes on the Earth's surface because its light is swamped in the glare of its brighter companion. Astronomers have long relied on a fundamental theoretical relationship between the mass of a white dwarf and its diameter. The theory predicts that the more massive a white dwarf, the smaller its diameter. The precise measurement of Sirius B's gravitational redshift allows an important observational test of this key



relationship.

The Hubble observations have also refined the measurement of Sirius B's surface temperature to be 25,000 degrees C. Sirius itself has a surface temperature of 10,000 degrees C.

At 8.6 light-years away, Sirius is one of the nearest known stars to Earth. Stargazers have watched Sirius since antiquity. Its diminutive companion, however, was not discovered until 1862, when it was first glimpsed by astronomers examining Sirius through one of the most powerful telescopes of that time.

Source: ESA/Hubble Information Centre

Citation: Hubble 'weighs' Dog Star's companion (2005, December 13) retrieved 5 May 2024 from <u>https://phys.org/news/2005-12-hubble-dog-star-companion.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.