

Astronomers Report Unprecedented Double Helix Nebula Near Center of the Milky Way

March 16 2006



The double helix nebula. (The image uses false colors because the eye is not sensitive to infrared light.) The spots are infrared-luminous stars, mostly red giants and red supergiants. Many other stars are present in this region, but are too dim to appear even in this sensitive infrared image. Credit: NASA/JPL-Caltech/UCLA

Astronomers report an unprecedented elongated double helix nebula near the center of our Milky Way galaxy, using observations from NASA's Spitzer Space Telescope. The part of the nebula the astronomers observed stretches 80 light years in length. The research is published

March 16 in the journal *Nature*.

"We see two intertwining strands wrapped around each other as in a DNA molecule," said Mark Morris, a UCLA professor of physics and astronomy, and lead author. "Nobody has ever seen anything like that before in the cosmic realm. Most nebulae are either spiral galaxies full of stars or formless amorphous conglomerations of dust and gas — space weather. What we see indicates a high degree of order."

The double helix nebula is approximately 300 light years from the enormous black hole at the center of the Milky Way. (The Earth is more than 25,000 light years from the black hole at the galactic center.)

The Spitzer Space Telescope, an infrared telescope, is imaging the sky at unprecedented sensitivity and resolution; Spitzer's sensitivity and spatial resolution were required to see the double helix nebula clearly.

"We know the galactic center has a strong magnetic field that is highly ordered and that the magnetic field lines are oriented perpendicular to the plane of the galaxy," Morris said. "If you take these magnetic field lines and twist them at their base, that sends what is called a torsional wave up the magnetic field lines.

"You can regard these magnetic field lines as akin to a taut rubber band," Morris added. "If you twist one end, the twist will travel up the rubber band."

Offering another analogy, he said the wave is like what you see if you take a long loose rope attached at its far end, throw a loop, and watch the loop travel down the rope.

"That's what is being sent down the magnetic field lines of our galaxy," Morris said. "We see this twisting torsional wave propagating out. We

don't see it move because it takes 100,000 years to move from where we think it was launched to where we now see it, but it's moving fast — about 1,000 kilometers per second — because the magnetic field is so strong at the galactic center — about 1,000 times stronger than where we are in the galaxy's suburbs."

A strong, large-scale magnetic field can affect the galactic orbits of molecular clouds by exerting a drag on them. It can inhibit star formation, and can guide a wind of cosmic rays away from the central region; understanding this strong magnetic field is important for understanding quasars and violent phenomena in a galactic nucleus. Morris will continue to probe the magnetic field at the galactic center in future research.

This magnetic field is strong enough to cause activity that does not occur elsewhere in the galaxy; the magnetic energy near the galactic center is capable of altering the activity of our galactic nucleus and by analogy the nuclei of many galaxies, including quasars, which are among the most luminous objects in the universe. All galaxies that have a well-concentrated galactic center may also have a strong magnetic field at their center, Morris said, but so far, ours is the only galaxy where the view is good enough to study it.

Morris has argued for many years that the magnetic field at the galactic center is extremely strong; the research published in Nature strongly supports that view.

The magnetic field at the galactic center, though 1,000 times weaker than the magnetic field on the sun, occupies such a large volume that it has vastly more energy than the magnetic field on the sun. It has the energy equivalent of 1,000 supernovae.

What launches the wave, twisting the magnetic field lines near the center

of the Milky Way? Morris thinks the answer is not the monstrous black hole at the galactic center, at least not directly.

Orbiting the black hole like the rings of Saturn, several light years away, is a massive disk of gas called the circumnuclear disk; Morris hypothesizes that the magnetic field lines are anchored in this disk. The disk orbits the black hole approximately once every 10,000 years.

"Once every 10,000 years is exactly what we need to explain the twisting of the magnetic field lines that we see in the double helix nebula," Morris said.

Co-authors on the *Nature* paper are Keven Uchida, a former UCLA graduate student and former member of Cornell University's Center for Radiophysics and Space Research; and Tuan Do, a UCLA astronomy graduate student. Morris and his UCLA colleagues study the galactic center at all wavelengths.

Source: UCLA

Citation: Astronomers Report Unprecedented Double Helix Nebula Near Center of the Milky Way (2006, March 16) retrieved 25 April 2024 from <https://phys.org/news/2006-03-astronomers-unprecedented-helix-nebula-center.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.