

Tibet Pathway for Chemicals To Reach Stratosphere

May 10 2006



A trio of NASA satellites observe in synchrony the vertical structures of thunderstorms (lower track) and their influences on ice clouds (color shades), water vapor (contours) and pollutants just above Earth´s lower atmosphere (higher track). Image courtesy of Rong Fu, Cinda Gillilan, Jonathan H. Jiang and Brian Knosp.

NASA and Georgia Tech researchers have found that thunderstorms over Tibet provide a main pathway for water vapor and chemicals to travel from the lower atmosphere, where human activity directly affects



atmospheric composition, into the stratosphere, where the protective ozone layer resides.

Learning how water vapor reaches the stratosphere can help improve climate prediction models. Similarly, understanding the pathways that ozone-depleting chemicals can take to reach the stratosphere is essential for understanding future threats to the ozone layer, which shields Earth from the sun's harmful ultraviolet rays.

Researchers from the Georgia Institute of Technology, Atlanta; NASA's Jet Propulsion Laboratory, Pasadena, Calif.; and the University of Edinburgh, Scotland, performed their analysis using data from the Microwave Limb Sounder instrument on NASA's Aura spacecraft, combined with data from NASA's Aqua and Tropical Rainfall Measuring Missions.

The team collected more than 1,000 measurements of high concentrations of water vapor in the stratosphere over the Tibetan Plateau and the Asian monsoon region. The measurements were collected during August 2004 and August 2005, during the height of monsoon season. Through the use of wind data and NASA atmospheric models, they found the water vapor originated over Tibet, just north of the Himalayan mountain range.

The team also found that even though more thunderstorms occurred over India, the storms over Tibet transported nearly three times more water vapor into the lower stratosphere than the more frequent thunderstorms that occur over India.

"This study shows that thunderstorms over Tibet are mainly responsible for the large amount of water vapor entering the stratosphere," said Dr. Rong Fu, associate professor in Georgia Tech's School of Earth and Atmospheric Sciences, who led the study. "The rainfall may not be as



frequent over Tibet as over the Indian monsoon area, but because Tibet is at a much higher elevation than India, the storms over Tibet are strong and penetrate very high, and send water vapor right into the stratosphere."

The study also found that the same pathway is responsible for transporting carbon monoxide, an indicator of air pollution, into the upper atmosphere.

"There's almost no carbon monoxide production in Tibet, so it's widely believed that carbon monoxide is transported to the tropopause over Southeast Asia and the Indian subcontinent," Fu said. The tropopause divides the lower atmosphere from the stratosphere, and is located at an altitude of about 18 kilometers (11 miles) above Earth over the tropics and Tibet.

Fu added, "Our study finds thunderstorms over Tibet transport as much carbon monoxide to the lower stratosphere as do those over India. When long-lived pollutants are transported out of the lower atmosphere, they can move rapidly. Pollutants from Asia, for example, can wind up on the other side of the world."

The findings are published in the *Proceedings of the National Academy of Sciences*.

Source: Georgia Institute of Technology

Citation: Tibet Pathway for Chemicals To Reach Stratosphere (2006, May 10) retrieved 16 June 2024 from <u>https://phys.org/news/2006-05-tibet-pathway-chemicals-stratosphere.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.