

Antarctic ocean found crucial to atmosphere's health

July 12 2006

Circulation in the waters near the Antarctic coast may be one of the planet's critical means of regulating levels of carbon dioxide in the Earth's atmosphere, according to researchers from MIT, Princeton and the National Oceanic and Atmospheric Administration.

Though climate scientists have long debated why atmospheric levels of carbon dioxide vary over lengthy periods in Earth's history, researchers now appear to have found a clue.

In a recent issue of the journal *Nature*, the team reports that computer modeling has revealed that the waters in the Southern Ocean below 60 degrees south latitude -- the region that hugs the continent of Antarctica -- play a far more significant role than was previously thought in regulating atmospheric carbon.

The waters north of this region do comparably little to regulate it, confuting past theories, the team found.

"Cold water that wells up regularly from the depths of the Southern Ocean spreads out on the ocean's surface along both sides of this dividing line, and we have found that the water performs two very different functions depending on which side of the line it flows toward," said Irina Marinov, the study's lead author and a postdoctoral fellow in MIT's Department of Earth, Atmospheric and Planetary Sciences.

"While the water north of the line generally spreads nutrients throughout

the world's oceans, the second, southward-flowing stream soaks up carbon dioxide, a greenhouse gas, from the air. Such a sharply defined difference in function has surprised us. It could mean that a change to one side of the cycle might not affect the other as much as we once suspected."

The research team also includes Professor Jorge Sarmiento of Princeton as well as Anand Gnanadesikan and Robbie Toggweiler of the National Oceanic and Atmospheric Administration (NOAA).

Two years ago, Sarmiento and colleagues discovered that the nutrients in the world's oceans were dependent on the Southern Ocean's circulation pattern. Scientists have also been aware that cold Antarctic waters have the ability to absorb atmospheric carbon dioxide.

What has been difficult is drawing the distinctions between one effect and another.

"The new paper shows that carbon dioxide and nutrient flow are separated quite dramatically," Sarmiento said. "What we are trying to do is understand better the balance of forces that help our planet maintain a steady environmental state, so we can anticipate what might cause that state to change. This paper helps us clarify how those forces interact."

Changing levels of atmospheric carbon dioxide have long concerned the scientific community, as this well-known greenhouse gas could be a major influence on global warming. Marinov said the discovery could shed light on how the Earth reacted far back in history, which might offer clues to how it will behave in the future.

"In the last ice age, for example, the atmosphere experienced very low levels of carbon dioxide, and no one is completely sure why," she said. "However, we now understand the Southern Ocean plays a large role in

regulating how much of the gas gets dissolved in water, and how much remains in the atmosphere."

The current study, she said, indicates that to better understand the Southern Ocean's effect on atmospheric carbon, scientists should pay greater attention to the Antarctic than to the more northerly sub-Antarctic region.

"In the Antarctic, the circulation pattern moves the surface water carrying carbon dioxide deep into the ocean's depths, where the sequestered carbon could potentially be trapped for a long time," Marinov said. "According to the models we used, the deep Antarctic is the critical region where we need to concentrate our research."

The team also indicated that the findings had implications for future research into carbon sequestration, a strategy for coping with increased atmospheric carbon dioxide levels. Some scientists propose that sequestration could one day capture atmospheric carbon and store it in places such as the deep ocean, thus mitigating humanity's greenhouse gas emissions.

"An interesting idea of recent years is that we can sequester a lot of carbon if we dump iron into the ocean to encourage the growth of certain microorganisms, which incorporate carbon as they grow," Marinov said. "These organisms would then fall to the ocean floor after they die, taking the carbon with them. The overall effect would be to lower the concentration of carbon in the surface waters, allowing more atmospheric carbon dioxide to dissolve into the sea. Our research has implications for future iron fertilization experiments, the focus of which we conclude should shift to the Antarctic."

Marinov said that the findings were based strongly on the team's computer models, which have limitations that they will now concentrate

on eliminating.

"While we are confident about the paper's conclusions, we are always looking for ways to clarify our understanding of the Southern Ocean," she said. "Our model, for example, does not take into account the fact that the circulation patterns are strongest in the winter, when the Antarctic is covered in darkness and the phytoplankton cannot grow very much. It is important that we understand the impact of this process on atmospheric carbon dioxide through future research."

Source: MIT

Citation: Antarctic ocean found crucial to atmosphere's health (2006, July 12) retrieved 19 April 2024 from <https://phys.org/news/2006-07-antarctic-ocean-crucial-atmosphere-health.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.