

## Solitons could power molecular electronics, artificial muscles

July 6 2006



A solitary electron wave (ripples of red and blue, center) travels along a polymer chain, causing the chain to bend in the middle. Image courtesy of Ohio State University.

Scientists have discovered something new about exotic particles called solitons. Since the 1980s, scientists have known that solitons can carry an electrical charge when traveling through certain organic polymers. A new study now suggests that solitons have intricate internal structures.

Scientists may one day use this information to put the particles to work



in molecular electronics and artificial muscles, said Ju Li, assistant professor of materials science and engineering at Ohio State University.

Li explained that each soliton is made up of an electron surrounded by other particles called phonons. Just as a photon is a particle of light energy, a phonon is a particle of vibrational energy.

The new study suggests that the electron inside a soliton can attain different energy states, just like the electron in a hydrogen atom.

"While we know that such internal electronic structures exist in all atoms, this is the first time anyone has shown that such structures exist in a soliton," Li said.

The soliton's quantum mechanical properties -- including these newly discovered energy states -- are important because they affect how the particle carries a charge through organic materials such as conducting polymers at the molecular level.

"These extra electronic states will have an effect -- we just don't know right now if it will be for better or worse," he said.

Li and his longtime collaborators from MIT published their findings in a recent issue of the *Proceedings of the National Academy of Sciences* (PNAS).

The name "soliton" is short for "solitary wave." Though scientists often treat particles such as electrons as waves, soliton waves are different. Ordinary electron waves spread out and diminish over time, and soliton waves don't.

"It's like when you make a ripple in water -- it quickly spreads and disappears," Li said. "But a soliton is a strange kind of object. Once it is



made, it maintains its character for a long time."

In fiber optics, normal light waves gradually flatten out; unless the signal is boosted periodically, it disappears. In contrast, solitonic light waves retain their structure and keep going without assistance. Some telecommunication companies have exploited that fact by using solitons to cheaply send signals over long distances.

Before solitons can be fully exploited in a wider range of applications, scientists must learn more about their basic properties, Li said. He's especially interested in how solitons carry a charge through conducting polymers, which consist of long, skinny chains of molecules.

The tiny chains are practically one-dimensional, and this calls some strange physics into play, Li said.

In their PNAS paper, Li and MIT colleagues Xi Lin, Clemens Furst, and Sidney Yip describe a detailed calculation of what happens to solitons at a quantum-mechanical level as they travel along a chain of the organic polymer polyacetylene.

Their mathematical model builds upon a 1979 model called the Su-Schrieffer-Heeger (SSH) model. Alan Heeger, a University of California, Santa Barbara physicist who discovered solitons, won the Nobel Prize in 2000 for his pioneering work on conducting polymers.

Li said the new work extends the SSH model by including the full flexibility of the polymer chain, as well as interactions between electrons.

The finding will likely affect the development of molecular electronics -- devices built from individual molecules.



Because polymer chains tend to bend and twist as solitons pass through them, scientists have wondered whether solitons could be used to power artificial muscles for high-tech robots and devices to aid human mobility. Such muscles would be made of organic polymers, and flex in response to light or electrochemical stimulation.

"If fully understood, solitons may also be harnessed to drive molecular motors in nanotechnology," Li said.

Source: by Pam Frost Gorder, Ohio State University

Citation: Solitons could power molecular electronics, artificial muscles (2006, July 6) retrieved 28 April 2024 from https://phys.org/news/2006-07-solitons-power-molecular-electronics-artificial.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.