


Hubble Probes Layer-Cake Structure of Alien World's Atmosphere

January 31 2007




Puffed-Up Atmosphere of a Star-Hugging Gas Giant Planet - This artist's illustration shows an extrasolar planet orbiting very close to its host star. The planet, designated HD 209458b, is about the size of Jupiter. Unlike Jupiter, the planet is so hot that its atmosphere is 'puffed up.' Starlight is heating the planet's atmosphere, causing hot gas to escape into space like steam rising from a boiler. HD 209458b completes an orbit every 3.5 days. The Hubble Space Telescope could not image the planet directly because it is too close to the star. Astronomers used Hubble to analyze the starlight that filtered through the planet's atmosphere. Imprinted on the starlight is information about the atmosphere's structure and chemical makeup. (Credit: NASA, ESA and G. Bacon of STScI)

The powerful vision of NASA's Hubble Space Telescope has allowed

astronomers to study for the first time the layer-cake structure of the atmosphere of a planet orbiting another star. Hubble discovered a dense upper layer of hot hydrogen gas where the super-hot planet's atmosphere is bleeding off into space.

The planet, designated HD 209458b, is unlike any world in our solar system. It orbits so close to its star and gets so hot that its gas is streaming into space, making the planet appear to have a comet-like tail. This new research reveals the layer in the planet's upper atmosphere where the gas becomes so heated it escapes like steam rising from a boiler.

“The layer we studied is actually a transition zone where the temperature skyrockets from about 1,340 degrees Fahrenheit (1,000 Kelvin) to about 25,540 degrees (15,000 Kelvin), which is hotter than the Sun,” said Gilda Ballester of the University of Arizona in Tucson, leader of the research team. “With this detection we see the details of how a planet loses its atmosphere.”



The solar-type star, HD 209458, and its 'hot Jupiter' planet in transit are shown in simulated violet light. As in this illustration, the star would appear as a limb-

darkened purple disk if seen in near-ultraviolet and violet light. The newly detected dense, narrow layer of hot hydrogen atoms is represented by the dark absorbing ring around the opaque planetary disk. The bulk of hydrogen atoms in the upper atmosphere, which forms an extensive cloud and also a comet-like tail, is shown in white. The absorbing layer was drawn at twice the altitude and 10 times the thickness to be more easily visible in this illustration, the rest of which has been drawn to scale. (Art credit: Loretta McKibben, UA Lunar and Planetary Lab).

The findings by Ballester, David K. Sing, of the University of Arizona and the Institut d'Astrophysique de Paris, and Floyd Herbert of the University of Arizona will appear Feb. 1 in a letter to the journal *Nature*.

Intense ultraviolet radiation from the host star heats the gas in the upper atmosphere, inflating the atmosphere like a balloon. The gas is so hot that it moves very fast and escapes the planet's gravitational pull at a rate of 10,000 tons a second, more than three times the rate of water flowing over Niagara Falls. The planet, however, will not wither away any time soon. Astronomers estimate its lifetime is more than 5 billion years.

The scorched planet is a big puffy version of Jupiter. In fact, it is called a "hot Jupiter," a large gaseous planet orbiting very close to its parent star. Jupiter might even look like HD 209458b if it were close to the Sun, Ballester said.

The planet completes an orbit around its star every 3.5 days. It orbits 4.7 million miles from its host, 20 times closer than the Earth is to the Sun. By comparison, Mercury, the closest planet to our Sun, is 10 times farther away from the Sun than HD 209458b is from its star. Unlike HD 209458b, Mercury is a small ball of iron with a rocky crust.

“This planet’s extreme atmosphere could yield insights into the atmospheres of other hot Jupiters,” Ballester said.

Although HD 209458b does not have a twin in our solar system, it has plenty of relatives beyond our solar system. About 10 to 15 percent of the more than 200 known extrasolar planets are hot Jupiters. A recent Hubble survey netted 16 hot Jupiter candidates in the central region of our Milky Way Galaxy, suggesting that there may be billions of these gas-giant star huggers in our galaxy.

HD 209458b is one of the most intensely studied extrasolar planets because it is one of the few known alien worlds that can be seen passing in front of, or transiting, its star, causing the star to dim slightly. In fact, the gas giant is the first such alien world discovered to transit its star. HD 209458b is 150 light-years from Earth in the constellation Pegasus.

The transits allow astronomers to analyze the structure and chemical makeup of the gas giant’s atmosphere by sampling the starlight that passes through the planet’s atmosphere. The effect is similar to finding fingerprints on a window by watching how sunlight filters through the glass.

Previous Hubble observations revealed oxygen, carbon, and sodium in the planet’s atmosphere, as well as a huge hydrogen upper atmosphere with a comet-like tail. These landmark studies provided the first detection of the chemical makeup of an extrasolar planet’s atmosphere.

Additional observations by NASA’s Spitzer Space Telescope captured the infrared glow from the planet’s hot atmosphere.

The new study by Ballester and her team is based on an analysis of archival observations made in 2003 with Hubble’s Space Telescope Imaging Spectrograph by David Charbonneau of the Harvard-

Smithsonian Center for Astrophysics in Cambridge, Mass. Ballester's team analyzed spectra from hot hydrogen atoms in the planet's upper atmosphere, a region not studied by Charbonneau's group.

Source: By Donna Weaver/Ray Villard, Space Telescope Science Institute

Citation: Hubble Probes Layer-Cake Structure of Alien World's Atmosphere (2007, January 31) retrieved 1 February 2023 from

<https://phys.org/news/2007-01-hubble-probes-layer-cake-alien-world.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.