


'Speechless' and 'Mute' help break the silence of the leaves

January 18 2007



The gene Speechless drives the division that initiates stomatal development. (Without Speechless, cells become the wax-coated, interlocking pavement cells on the surface of the plant, none of which open or close.) Two or three divisions after Speechless, Mute causes the formation of a guard mother cell. Then Fama causes the guard mother cell to divide into two equal guard cells that, as their name implies, guard the opening of the stomatal pore. Credit: University of Washington

Researchers have discovered two genes that guide land plants to develop microscopic pores that they can open and close as if each pore was a tiny mouth.

Plants wouldn't have been able to move from water to land 400 million years ago if they hadn't evolved this ability, which protects them from losing too much moisture.

The leaves and stems of land plants are dotted with the "tiny mouths," called stomata. When open, stomata allow the plant to take in carbon dioxide gas needed for photosynthesis and allow moisture to evaporate, pulling water from the roots into the plant. But when too much moisture is being lost, the two cells around the stomatal pore close it completely.

Without the genes guiding stomatal development, plants won't develop any mouthlike pores, hence the names Speechless and Mute for the newly discovered genes, according to Keiko Torii, a University of Washington associate professor of biology.

Two separate papers on the genes, one by Torii's UW group and the other by Stanford University researchers, have been published online by *Nature*, and are scheduled to appear in the print publication Feb. 1. Each group describes independently finding the gene that came to be called Speechless and its role in initiating the process that leads to stomata.

In addition, Torii's UW group published findings in its *Nature* article about another gene, one they named Mute, that triggers the key middle step that decides when a cell will fully become a stomata. Earlier this year the Stanford group published findings about the gene that controls the final step in stomata development, called Fama.

"In the last few months, we've gone from knowing surprisingly little about the genes involved to knowing all three major factors – Speechless, Mute and Fama," says Lynn Pillitteri, a research associate in biology and lead author of the *Nature* paper.

That the three are so closely related will be of interest to biologists studying both plants and animals, she says. Each is a basic protein with a helix-loop-helix domain, a sequence that is quite ancient and controls a vast range of physiological and developmental processes. Speechless, Mute and Fama also have very similar DNA sequences and could have

arisen from a single gene that replicated and evolved, giving plants additional genes with slightly different characteristics.

Having two or three genes with similar characteristics would give plants what Torii terms "the freedom to play, to make functions that are the more elaborate stomata in modern plants."

Other biologists have seen something similar in animals. The ability to differentiate cells that become muscles also is controlled by consecutive action of basic helix-loop-helix proteins with DNA closely related to each other.

Molecular conservation of such key regulatory genes between plants and animals – genes that switch on and off cell-type differentiation programs from precursor stem cells – is intriguing and exciting, Torii says.

Source: University of Washington

Citation: 'Speechless' and 'Mute' help break the silence of the leaves (2007, January 18) retrieved 26 April 2024 from <https://phys.org/news/2007-01-speechless-mute-silence.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.