

# Lakes beneath Antarctic ice sheets found to initiate and sustain flow of ice to ocean

February 21 2007

---

The Earth Institute at Columbia University—One of the planet's most remote and little-understood features may play a crucial role in transporting ice from the remote interior of Antarctica towards the surrounding ocean according to a new research.

Geophysicists Robin Bell and Michael Studinger from the Lamont-Doherty Earth Observatory, a part of The Earth Institute at Columbia University, led a team that discovered four large, subglacial lakes that for the first time link these water bodies locked beneath miles of ice, to fast flowing ice streams in Antarctica. Together with colleagues from NASA, the University of New Hampshire and the University of Washington, the scientists found that, in four separate cases, lakes appear to contribute to the formation of ice streams. Their work appears in the February 22 issue of the journal *Nature*.

Ice streams are large, fast-flowing features within ice sheets that transport land-based ice and meltwater to the ocean. One such stream, the Recovery Ice Stream, drains 8 percent of the U.S.-sized East Antarctic Ice Sheet. The Recovery basin, unexplored since 1966, funnels an estimated 35 billion tons of ice into the Weddell Sea annually.

"Until about a year ago, not many people cared much about subglacial lakes," said Studinger. "That's changing, but we're still only just beginning to understand how these lakes, sealed beneath more than two miles of ice, have the potential to impact the rest of the world."

The scientists examined satellite radar images and high-resolution laser profiles of the region for ice stream patterns and surface features indicating the presence of subglacial lakes beneath the ice. Not only did they find four new lakes, but they discovered that the lakes coincide with the origin of tributaries of the Recovery Glacier. Upstream of the lakes, the ice sheet moves at just 2 to 3 meters per year; downstream the flow increases to nearly 50 meters per year. Bell and Studinger conclude that the lakes provide a reservoir of water that lubricates the bed of the stream to facilitate ice flow and prevent the base of the sheet from freezing to the bedrock.

Moreover, their work suggests that subglacial lakes could play a role in and sea level rise as well as regional and global climate change. Meltwater at the base of ice streams increases the flow of ice to the oceans, which could, in turn, contribute to higher sea levels worldwide. In addition, floods have been known to originate from the interior of the ice sheet in the past, possibly from subglacial lakes. These sudden pulses of fresh water could potentially interfere with nearby ocean currents that redistribute heat and carbon dioxide around the globe, disrupting the Earth's finely tuned climate system.

"It's almost as if the lakes are capturing the geothermal energy from the entire basin and releasing it to the ice stream." said Bell. "They power the engines that drive ice sheet collapse. The more we learn about them, the more we realize how important they are."

Source: The Earth Institute at Columbia University

Citation: Lakes beneath Antarctic ice sheets found to initiate and sustain flow of ice to ocean (2007, February 21) retrieved 25 April 2024 from <https://phys.org/news/2007-02-lakes-beneath-antarctic-ice-sheets.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.