

Long-lived deep-sea fishes imperiled by technology, overfishing

February 19 2007

Many commercially prized fish from the depths of the world's oceans are severely threatened by over-fishing and the species' ability to recover is constrained by the fishes' long lifespans and low reproductive success, a panel of experts said today at the annual meeting of the American Association of the Advancement of Science. Some of the fish species living at depths greater than 500 meters take decades to reach breeding maturity, so there are no quick-fix remedies available to replenish the population, said Selina Heppell, a fisheries biologist from Oregon State University.

"The harvest of deep-sea fishes is a lot like the harvest of old-growth timber," Heppell said, "except we don't 'replant' the fish. We have to depend on the fish to replenish themselves. And the habitat that used to provide them protection – the deep ocean – is now accessible to fishing because of new technologies."

Among the most recognized deep-sea species at-risk are orange roughy and Patagonian toothfish, better known as Chilean sea bass. In the deep ocean off the Pacific Northwest, sablefish – also known as black cod – are another depleted species. Deep-sea fishes grow slowly because of limited food sources and slower metabolisms and many don't reach sexual maturity for 30 to 40 years, Heppell said. The harvest of older fish may have an even greater impact on these threatened populations because older fish are more likely to breed successfully than younger fish.

"When you buy orange roughy at the store, you are probably purchasing a filet from a fish that is at least 50 years old," Heppell said. "Most people don't think of the implications of that. Perhaps we need a guideline that says we shouldn't eat fish that are as old as our grandmothers."

Most of the deep-sea fishes are in international waters, where there are no guidelines and protections – unlike within United States territorial waters. Most of these fish are caught by deep trawlers near seamounts, where they congregate because of food.

Technological advances have made targeting these fish easier, the panelists pointed out, because powerful ships can drag huge nets hundreds of feet below the surface. New refrigeration techniques, including "flash freezing," allow ships to range far out into the ocean for days at a time. And sophisticated global positioning systems (GPS) and fish finders can target schools of fish or seamounts with ease.

"One reason that many of these fish species were fished sustainably in the past is that we couldn't fish all of the places all of the time," Heppell said. "That isn't necessarily true anymore."

Heppell is a faculty member in the Department of Fisheries and Wildlife at Oregon State, where she studies fish dynamics, populations and life cycles. The deep-sea fishes are among the hardest to study for obvious reasons, she said, and additional research is critical to protect these species.

Long-lived fish usually have low reproductive rates, either because of low breeding success or high mortality. In the case of deep-sea fishes, both scenarios often play out.

In some species, such as sharks, the fish may only produce a handful of

offspring and the chances of survival by an individual are low. In other species, including orange roughy and oreos, an individual fish produces thousands of eggs – most of which die through predation or starvation.

"One of the things we need to know more about is how the fish larvae get transported," Heppell said. "We don't know whether fish from different seamounts are genetically distinct or whether larvae from one seamount end up populating another. The odds against these fish are so high that, in a reproductive sense, they have to wait for the stars to align before they successfully produce offspring that will survive until maturity."

Natural and human-influenced climate factors including El Niño, the Pacific Decadal Oscillation and hypoxia zones all can influence shallow-water fishes' breeding and mortality rates, but deep-sea environments are usually stable. The deep sea is almost completely dark, very near freezing and has very little food – reasons for the fishes' slow growth and low productivity.

"Old fish don't necessarily need to breed every year," Heppell pointed out, "so when nature throws a bad reproductive year at them, the species can survive. But the point remains that you have to have older fish to replenish the stock when those bad years come."

Conversely, Heppell said, good years often can carry the population in a phenomenon known as "episodic recruitment." In studies of long-lived fish species, it isn't unusual to find a school with numerous 18-year-old fish, for example, but very few fish that are 17 or 19 years of age. Scientists can determine the age of fish through their otoliths, or ear bones, which regularly accumulate rings much like trees. Variations in the size of the rings can indicate ocean productivity that year.

Harvesting older fish lessens the likelihood of many productive breeding

years, Heppell pointed out, and lengthens the time species need to recover.

"There are models that estimate the recovery time for some rockfish species is at least 200 years," Heppell said. "And we still don't know all of the factors that influence their survival."

Source: Oregon State University

Citation: Long-lived deep-sea fishes imperiled by technology, overfishing (2007, February 19) retrieved 27 April 2024 from <https://phys.org/news/2007-02-long-lived-deep-sea-fishes-imperiled-technology.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.