

Ocean iron and CO₂ interaction studied

April 26 2007

A French study suggested that iron supply changes from deep water to the ocean's surface might have a greater effect on atmospheric CO₂ than thought.

The finding by Stephane Blain and colleagues at the University of the Mediterranean have implications for interpretations of past climate change, as well as future climate predictions.

Iron plays an important role in the carbon cycle because phytoplankton need iron to fix energy into carbon, which is then sequestered in the deep ocean when the microorganisms die and sink. But the exact effect of changes in iron supply on the amount of carbon sequestered remains unclear.

The researchers studied a phytoplankton bloom that occurs in the Southern Ocean and found the ratio of carbon exported to the ocean interior to the amount of iron supplied is at least 10 times higher than previous thought.

The difference occurs because the iron is supplied slowly and continuously and because the bloom is dependent on the supply of major nutrients from below. That indicates such high efficiencies are unlikely to be achieved through artificial "fertilization" of the ocean with iron.

The research appears in the journal *Nature*.

Copyright 2007 by United Press International

Citation: Ocean iron and CO₂ interaction studied (2007, April 26) retrieved 28 April 2024 from <https://phys.org/news/2007-04-ocean-iron-co2-interaction.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.