


Bat flight generates complex aerodynamic tracks

May 10 2007



On the cover of *Science*: A bat, *Glossophaga soricina*, in flight in a wind tunnel at Lund University, Sweden. The velocity field induced by the wingbeat is shown superimposed and to scale. The bat wake reconstructions have features that have not been observed in similar-sized birds. Membranous and feathered wings of natural and engineered flyers may have significantly different aerodynamic properties. Credit: Lund University, Sweden

Bats generate a measurably distinct aerodynamic footprint to achieve lift and maneuverability, quite unlike birds and contrary to many of the assumptions that aerodynamicists have used to model animal flight,

according to University of Southern California aerospace engineer Geoffrey Spedding.

The researcher, together with a multi-institutional team of scientists found that bat flight is quite different from bird flight, particularly at very small scales. They based their findings on new measurements of aerodynamic performance in the wing beats of a small species of bat.

"Bats with a body mass of 10 – 30 grams — or about the weight of one or two teaspoons of sugar — and tip-to-tip wing spans of 25 – 30 centimeters — about the length of a human hand — generate very different wakes," he reported in the May 11, 2007 issue of *Science*.

"The tell-tale tracks in the airflow caused by the wing beat have a very different pattern for bats, and this difference can be traced to the peculiar upstroke," he continued. "That, in turn, is likely caused by the collapsible membrane of the bat's wing, which needs to maintain some degree of tension."

His research group's findings are presented in an article titled "Bat Flight Generates Complex Aerodynamic Tracks." Coauthors included A. Hedenström, L. C. Johansson and M. Wolf of Lund University, Sweden; R. von Busse of the University of Munich, Germany; and Y. Winter at Bielefeld University in Germany.

Bats constitute about 20 percent of mammalian species, but not much attention has been paid to them because of their nocturnal habits. Most bats eat flying insects, consuming up to their body weight each night. Their agility and tremendous ability to maneuver in flight is accentuated by their ability to locate prey using an advanced echo-location system, which is basically a sonar operated by ultrasound.

The authors noted that bat wings are very different from those of birds

because of their separate evolutionary paths to flight.

"Instead of feathers projecting back from lightweight, fused arm and hand bones, bats have flexible, elastic membranes that stretch between specially extended, slender bones of the hand," Spedding said. "The bones and wing membrane both change shape with every wing beat, flexing in response to the balance between forces applied by the muscles and competing forces due to the air motion around them. "

In contrast with bird wings, the bat wing membrane must be kept under tension, otherwise it will flap uselessly, like a flag. As a consequence, Spedding said, there are limits to how much the wing can be folded during flight.

Spedding and his colleagues think that the unique aerodynamic wake signatures of bat wings are caused by different mechanical operations in the upstroke of the wingbeat.

"Where birds can feather their wings, opening the feathers like a Venetian blind, bats must do something different," Spedding said. "Hence, they have developed a twisting wing path that increases the lift during the upstroke."

This is the first time that this positive aerodynamic effect has been observed in wing flapping, he said, and it is consistent with previous speculation that was based on the bat's complicated wing motions alone.

Spedding said the findings are just the beginning of an exciting research program to make detailed measurements of bat wings and add to a growing body of data about bat flight.

"Bats are agile hunters, capable of plotting and executing complex maneuvers through cluttered environments," he said. "These are the

traits we'd like our unmanned air vehicles to have because there are so many complex rural and urban environments in which we could use them."

With the promise of human-engineered, micro-scale flight just around the corner, engineers may also be able to emulate bat wings, building simple mechanical wings of flexible, elastic material stretched over support rods, to improve the aeronautical performance of smaller airborne vehicles.

"Bats have relied on very flexible wings for 50 million years to propel and lift themselves into the sky," Spedding said. "We still have a lot more to learn about the aerodynamics of bat flight and how their wings allow them to maneuver through incredibly unsteady air flows and turbulent conditions."

For a brief video clip of a bat at a feeder, taken at Lund University, Sweden, visit: viterbi.usc.edu/images/video/bat_flying.wmv

Source: University of Southern California

Citation: Bat flight generates complex aerodynamic tracks (2007, May 10) retrieved 28 January 2023 from <https://phys.org/news/2007-05-flight-complex-aerodynamic-tracks.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.