

Philandering female felines forgo fidelity

May 31 2007

Cheetah study reveals many litters have more than one father

While promiscuity in the animal kingdom is generally a male thing, researchers for the Wildlife Conservation Society and the Zoological Society of London (ZSL) have found that, in cheetah society, it's the female with the wandering eye, as reported in a paper in the latest issue of Proceedings of the Royal Society B.

According to the study, researchers found that about 43 percent of cheetah litters with more than one cub were fathered by more than one male, revealing a mating system that deviates from those used by other carnivores, most of which consist of single or sibling males monopolizing many females.

"It seems that female cheetahs are highly promiscuous with no detectable mate fidelity between breeding seasons," said Dr. Sarah Durant, a researcher with the Wildlife Conservation Society/Tanzania Cheetah Conservation Program and project leader. "In fact, we now know many of the fathers exist outside the park and study area. This is a finding which gives us a critical insight into cheetah ecology and one that merits inclusion into future conservation efforts for the species."

The nine-year study, which was conducted in the Serengeti National Park of Tanzania, focused on determining the paternity of cheetahs through the use of fecal samples that were then subjected to genetic testing. By comparing individual genetic micro-satellites from 171 samples, researchers not only were able to determine if more than one father contributed to a litter, but also which of the known male cheetahs



in the study area had sired cubs. Surprisingly, few known males had, with only 21.3 percent being assigned paternity, although not all males in the study area had been sampled. At least some of these missing fathers undoubtedly resided outside the study area.

The findings corroborate established studies on ranging patterns of female and male cheetahs. Female cheetahs occupy a home range averaging some 833 square kilometers in size, whereas a male's territory averages 37 square kilometers. In many carnivore species, it's the males that have larger ranges that encompass several female ranges.

"The picture that emerges is that of wide-ranging females mating with a number of males, which could be an advantage in terms of helping to maintain genetic diversity," said ZSL's Dada Gottelli, the study's lead author.

Findings from the study are significant for conservation efforts for the species, which is threatened by a number of human-related causes. Further, cheetahs have always existed in low densities, due to competition from larger predators such as the lion and hyena. "Our study shows that more fathers are contributing to the next generation than we thought," added Durant. "This is good news for cheetahs, and a reminder that future conservation strategies need to encompass large landscapes if females are to continue to have access to many males."

In addition to helping save cheetahs in Africa, WCS is also studying the last remaining populations of Asia's cheetahs in Iran.

Source: Wildlife Conservation Society

Citation: Philandering female felines forgo fidelity (2007, May 31) retrieved 19 April 2024 from



https://phys.org/news/2007-05-philandering-female-felines-forgo-fidelity.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.