

Spitzer Searches for the Origins of Life

June 13 2007



This false-color composite from NASA's Spitzer Space Telescope and NASA's Chandra X-ray Observatory shows the remnant of one a supernova, or the explosive death of a star. Image credit: NASA/JPL- Caltech/Harvard-Smithsonian CfA

Astronomers suspect the early Earth was a very harsh place. Temperatures were extreme, and the planet was constantly bombarded by cosmic debris. Many scientists believe that life's starting materials, or building blocks, must have been very resilient to have survived this tumultuous environment.

Now, NASA's Spitzer Space Telescope has learned, for the first time, that organic molecules believed to be among life's building blocks, called polycyclic aromatic hydrocarbons, can survive another type of harsh setting, an explosion called a supernova. Supernovae are the violent deaths of the most massive stars. In death, these volatile objects blast



tons of energetic waves into the cosmos, destroying much of the dust surrounding them.

The fact that polycyclic aromatic hydrocarbons can survive a supernova indicates that they are incredibly tough - like cosmic cockroaches enduring a nuclear blast. Such durability might be further proof that these molecules are indeed among life's building blocks.

Achim Tappe of the Harvard-Smithsonian Center for Astrophysics, Cambridge, Mass., used Spitzer's infrared spectrograph instrument to detect abundant amounts of polycyclic aromatic hydrocarbons along the ridge of supernova remnant N132D. The remnant is located 163,000 light-years away in a neighboring galaxy called, the Large Magellanic Cloud.

"The fact that we see polycyclic aromatic hydrocarbons surviving this explosion illustrates their resilience," says Tappe.

These intriguing molecules are comprised of carbon and hydrogen atoms, and have been spotted inside comets, around star-forming regions and planet-forming disks. Since all life on Earth is carbon based, astronomers suspect that some of Earth's original carbon might have come from these molecules - possibly from comets that smacked into the young planet.

Astronomers say there is some evidence that a massive star exploded near our solar system as it was just beginning to form almost 5 billion years ago. If so, the polycyclic aromatic hydrocarbons that survived that blast might have helped seed life on our planet.

Tappe's paper was published in the December 10, 2006, issue of *Astrophysical Journal*.



Source: by Whitney Clavin, JPL/NASA

Citation: Spitzer Searches for the Origins of Life (2007, June 13) retrieved 2 May 2024 from https://phys.org/news/2007-06-spitzer-life.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.