


Arizona State scientists keep an eye on Martian dust storm

July 11 2007



The map shows the atmosphere's opacity at an infrared wavelength of 9 micrometers. The scale bar's values run from nearly clear (0.05) to roughly a two-thirds reduction in sunlight (0.40). Image credit: NASA/JPL/Arizona State University

Scientists at Arizona State University's Mars Space Flight Center are using the Thermal Emission Imaging System (THEMIS) on NASA's Mars Odyssey orbiter to monitor a large dust storm on the Red Planet. The instrument, a multi-wavelength camera sensitive to five visible wavelengths and 10 infrared ones, is providing Mars scientists and spacecraft controllers with global maps that track how much atmospheric dust is obscuring the planet.

The dust storm, which erupted during the last week of June 2007, is

affecting operations for all five spacecraft operating at Mars. The fleet includes two NASA rovers on the ground (Spirit and Opportunity), plus three orbiters, two of which belong to NASA (Mars Odyssey and Mars Reconnaissance Orbiter) and one to the European Space Agency (Mars Express).

Beginning in Mars' heavily cratered southern highlands, the dust storm took roughly a week to grow large enough to encircle the planet. Dust has now drifted into the northern hemisphere as well.

"This is the favorable time of the Martian year for dust storms," says Joshua Bandfield, research associate at the Mars Space Flight Facility. The facility is part of the School of Earth and Space Exploration in ASU's College of Liberal Arts and Sciences.

"It's summer in the southern hemisphere," he says, "That's when Mars lies closest to the sun and solar heating is greatest."

Bandfield adds, "We can watch weather fronts spreading and kicking up dust in a big way." He explains that as winds sweep dust into the atmosphere, the atmosphere becomes warmer. This adds to the storm's power, helping it to pick up more dust. But the process has a built-in limitation, he says. "When the dust becomes thick enough, it reflects more sunlight from the atmosphere, allowing the air near the surface to cool."

As seen from orbit, the dust storm has the effect of veiling surface features – or even concealing them completely, which hasn't happened yet in this event. "This storm isn't as big or severe as the one in 2001," Bandfield says. "THEMIS and other orbiters can still see the surface, despite the continuing dust activity."

From the ground, the dust in the air has cut the amount of sunlight

reaching the rovers' solar panels and reducing their electrical power. "If you were standing there, you'd see the sky looking tawny with haze," explains Bandfield. "The sun would appear as a sharp-edged disk, but the light level would be noticeably lower than what you would see under a totally clear sky."

Luckily, say scientists, summer is a time when the rovers can best survive under reduced power. If the storm had struck during local winter, the rovers might not get enough power during the day to stay alive through the cold Martian night.

How long will this storm last? No one knows for sure, but Bandfield notes its effects won't disappear as quickly as the storm erupted. "Mars," he says, "will remain dusty for at least a couple months more."

Source: Arizona State University

Citation: Arizona State scientists keep an eye on Martian dust storm (2007, July 11) retrieved 17 July 2024 from <https://phys.org/news/2007-07-arizona-state-scientists-eye-martian.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.