

## Early fire risk for mountains near Los Angeles

July 2 2007



Philip Dennison, a University of Utah geographer, measures burned vegetation after Southern California's 2003 Simi Fire. Dennison has devised a method to forecast the highest-risk period of large fires in the Santa Monica Mountains near Los Angeles. Credit: Philip Dennison, University of Utah

Researchers at the University of Utah and elsewhere have developed a new way to predict when vegetation dries to the point it is most vulnerable to large-scale fires in the Santa Monica Mountains near Los Angeles. And this year's forecast says the highest-risk fire period will begin July 13 – weeks earlier than usual.

Despite that, the new study also shows that unlike other areas of the western United States, global warming has not caused any apparent long-term trend toward early fire seasons in the Santa Monicas.



The scientists eventually hope to expand their unique fire-risk forecasting method to all of Southern California and beyond.

"We developed a way to predict when the time of highest fire danger begins in the Santa Monica Mountains, based on the amount of spring precipitation," says the study's principal author, Philip Dennison, an assistant professor of geography at the University of Utah. "We estimate that this year, the highest fire danger will begin July 13."

The study found the amount of March-April-May precipitation can be used to predict the date at which high fire-risk thresholds are reached.

Dennison says he hopes that "in future years this method can be used to better plan for the start of high fire danger. Fire agencies could use this to help them plan where and when to put their people and equipment. Homeowners may find this useful for knowing when fire danger will be high so they can be better prepared to evacuate, clear brush from around their homes and watch for arsonists."

Study co-author Max Moritz, a wildland fire specialist at the University of California, Berkeley, adds: "Although large fires still may be associated with autumn Santa Ana winds in Southern California, it appears the stage for these events is set much earlier than usual this year. Being prepared early is thus even more important than usual."

The study will be published later this year in the International Journal of Wildland Fire. Dennison and Moritz conducted it with Robert S. Taylor, a fire specialist at the National Park Service's Santa Monica Mountains National Recreation Area.

## Narrowing the Start of the High-Risk Fire Season

Dennison says he is releasing the findings now because the new method



indicated that the moisture level in chamise – the 3- to 12-foot-tall, primary evergreen shrub in the Santa Monicas' predominant chaparral ecosystem – will reach a critically low "live fuel moisture" level of 71 percent July 13, with a one- to two-week margin of error.

In most years, such vulnerability to ignition is not reached until later. The scientists found that during the 1984-2005 study period, plant moisture dropped below 71 percent as early as June 22 and as late as Nov. 4. It usually happens in late August. In September, fierce Santa Ana winds become more likely to blow over Southern California.

"I don't think it will be a surprise that things are dry this year" throughout the West, Dennison says. "But if you can pin it to a short time period of a few weeks within a four-month time frame, then you are providing a useful prediction."

"You can focus your efforts on the period of highest fire danger. The study shows that the period of high fire danger is predictable months in advance, so by June 1, fire agencies can know the earliest date to expect large fires."

Dennison hopes to expand the forecast method to the Southern California region and nationally to other places where vegetation moisture is an important factor in fires.

Santa Anas are strong, hot, dry winds that blow over Southern California from the east or northeast, most frequently September through April but peaking in December.

The Santa Monica Mountains run west-to-east, paralleling the Pacific Coast near Malibu. The more urban eastern end includes the Hollywood Hills and Beverly Hills, which divide the Los Angeles Basin from the San Fernando Valley. The wilder western part of the range includes the



Santa Monica Mountains National Recreation Area.

The researchers call it "the world's largest urban national park" and "one of the major areas of wildland-urban interface and intermix in Southern California," located near Malibu, Calabasas, Thousand Oaks, Westlake Village, Newbury Park, Agoura Hills and parts of Los Angeles: Encino, Brentwood, Laurel Canyon and Woodland Hills.

## A History of Fire: The Study, Part 1

The study examined all 59 fires in the Santa Monicas during 1984 through 2005, and found that one of seven major fires – those burning more than 2,500 acres – occurred when chaparral vegetation moisture dropped to 77 percent. The other six big fires happened after conditions grew even drier and plant moisture had dropped to 71 percent.

The seven large fires account for 90 percent of the acreage burned in the Santa Monica Mountains from 1984 through 2005, Dennison says.

The researchers calculated that plant moisture falls to 71 percent on July 8 in years with no spring precipitation. They predicted that for each inch of spring rain, that moisture level will be reached 10 days later.

The researchers plan to refine their technique to pinpoint a single high fire-risk threshold between 77 percent and 71 percent foliage moisture. The uncertainty is caused by the one large fire that ignited when foliage moisture was 77 percent.

Dennison says researchers who conducted a 2006 study "found that fire season is starting earlier across the western United States, and they attributed it to earlier snowmelt which is caused by global warming."

Despite this year's predicted early fire season, the new study found no



trend toward an earlier start for fire seasons in the Santa Monicas during 1933-2005. Dennison says the Santa Monicas don't receive much snow, and it melts fast.

"Small fires can occur anytime" but large fires are most likely when plant moisture falls below the thresholds, he says. "Low live fuel moisture is one ingredient in producing large fires in Southern California. You also need high winds and something to start the fire."

The researchers tested their forecasting method back to 1933, and found the predicted date for vegetation moisture to drop below 71 percent was within one week of the predicted date for half of the years, and within two weeks for most years.

## Learning to Predict Dangerous Dryness: The Study, Part 2

The study also examined nine variables to determine which best predicted when vegetation moisture would drop to thresholds making the plants highly vulnerable to fire.

Factors that didn't predict plant moisture well included three variables that used U.S. weather satellite measurements of "greenness" in the Santa Monicas, and another three variables that measured moisture in plants before the fire season.

Three other variables involved rainfall amounts during various periods: the Nov. 1-May 31 wet season total, winter (December-January-February) precipitation and spring (March-April-May) precipitation.

The study found spring rainfall amounts at a University of California, Los Angeles, weather station can be used to make statistically valid



predictions of when "live fuel moisture" would drop to the 77 percent and 71 percent thresholds.

"Live fuel moisture" data came from the Los Angeles County Fire Department, which about every three weeks collects samples of chaparral vegetation at three locations in the western Santa Monicas: Clark Motorway, Schueren Road and Trippet Ranch.

The collected leaves and twigs are weighed, oven-dried and then weighed again. A "live fuel moisture" of 71 percent means the weight of water that was in the plant sample was 71 percent of the sample's dry weight.

Moritz says UC, Berkeley, has developed tools homeowners can use to prepare for fire season: <u>firecenter.berkeley.edu/toolkit/homeowners.html</u>

Source: University of Utah

Citation: Early fire risk for mountains near Los Angeles (2007, July 2) retrieved 1 May 2024 from <u>https://phys.org/news/2007-07-early-mountains-los-angeles.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.