


Interstellar Chemistry Gets More Complex With New Negatively-Charged Molecule Discovery

July 23 2007



Astronomers using the Robert C. Byrd Green Bank Telescope found the negatively-charged form of octatetraynyl (C_8H^-) in a cold interstellar cloud (middle left) and in the gaseous envelope surrounding an old, evolved star (middle right). This is the largest negatively-charged molecule yet found in space. The scientists believe it probably is formed in steps, illustrated here, proceeding downward. 1. A molecule of C_2H attaches to a molecule of C_6H_2 , producing a molecule of C_8H_2 and a hydrogen atom. 2. Radiation (squiggly line) breaks one hydrogen atom from the C_8H_2 , leaving C_8H and a hydrogen atom. 3. Finally, an electron attaches itself to the C_8H molecule, freeing a burst of radiation (overall glow seen around the molecule) and leaving the negatively-

charged ion C_8H^- . Credit: Bill Saxton, NRAO/AUI/NSF

Astronomers using data from the National Science Foundation's Robert C. Byrd Green Bank Telescope (GBT) have found the largest negatively-charged molecule yet seen in space. The discovery of the third negatively-charged molecule, called an anion, in less than a year and the size of the latest anion will force a drastic revision of theoretical models of interstellar chemistry, the astronomers say.

"This discovery continues to add to the diversity and complexity that is already seen in the chemistry of interstellar space," said Anthony J. Remijan of the National Radio Astronomy Observatory (NRAO). "It also adds to the number of paths available for making the complex organic molecules and other large molecular species that may be precursors to life in the giant clouds from which stars and planets are formed," he added.

A team of scientists from the Harvard-Smithsonian Center for Astrophysics (CfA) found negatively-charged octatetraynyl in a cold, dark cloud of molecular gas. A second team headed by Remijan found octatetraynyl in the envelope of gas around an old, evolved star. In both cases the molecule, a chain of eight carbon atoms and one hydrogen atom, had an extra electron, giving it a negative charge.

About 130 neutral and about a dozen positively-charged molecules have been discovered in space, but the first negatively-charged molecule was not discovered until late last year. The largest previously-discovered negative ion found in space has six carbon atoms and one hydrogen atom.

Ultraviolet light from stars can knock an electron off a molecule,

creating a positively-charged ion. Astronomers had thought that molecules would not be able to retain an extra electron, and thus a negative charge, in interstellar space for a significant time. “That obviously is not the case,” said Mike McCarthy of the CfA. “Anions are surprisingly abundant in these regions.”

“Until recently, many theoretical models of how chemical reactions evolve in interstellar space have largely neglected the presence of anions. This can no longer be the case, and this means that there are many more ways to build large organic molecules in cosmic environments than have been explored,” said Jan M. Hollis of NASA's Goddard Space Flight Center (GSFC).

Remijan and his colleagues found the octatetraynyl anions in the envelope of the evolved giant star IRC +10 216, about 550 light-years from Earth in the constellation Leo. They found radio waves emitted at specific frequencies characteristic of the charged molecule by searching archival data from the GBT, the largest fully-steerable radio telescope in the world.

Another team from the Harvard-Smithsonian Center for Astrophysics found the same characteristic emission when they observed a cold cloud of molecular gas called TMC-1 in the constellation Taurus. These observations also were done with the GBT. In both cases, preceding laboratory experiments by the CfA team showed which radio frequencies actually are emitted by the molecule, and thus told the astronomers what to look for.

“It is essential that likely interstellar molecule candidates are first studied in laboratory experiments so that the radio frequencies they can emit are known in advance of an astronomical observation,” said Frank Lovas of the National Institute of Standards and Technology (NIST).

Both teams announced their results in the July 20 edition of the *Astrophysical Journal Letters*.

“With three negatively-charged molecules now found in a short period of time, and in very different environments, it appears that many more probably exist. We believe that we can discover more new species using very sensitive and advanced radio telescopes such as the GBT, once they have been characterized in the laboratory,” said Sandra Bruenken of the CfA.

“Further detailed studies of anions, including astronomical observations, laboratory studies, and theoretical calculations, will allow us to use them to reveal new information about the physical and chemical processes going on in interstellar space,” said Martin Cordiner, of Queen's University in Belfast, Northern Ireland.

“The GBT continues to take a leading role in discovering, identifying and mapping the distribution of the largest molecules ever found in astronomical environments and will continue to do so for the next several decades,” said Phil Jewell of NRAO.

Source: Harvard-Smithsonian Center for Astrophysics

Citation: Interstellar Chemistry Gets More Complex With New Negatively-Charged Molecule Discovery (2007, July 23) retrieved 9 April 2024 from <https://phys.org/news/2007-07-interstellar-chemistry-complex-negatively-charged-molecule.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--