

# AKARI's observations of asteroid Itokawa

August 23 2007

---

The space-borne infrared observatory AKARI, observed asteroid Itokawa last month with its Infrared Camera. The data will be used to refine estimates of sizes of potentially hazardous asteroids in the future.

The data collected by AKARI, a JAXA mission with ESA participation, complements that from JAXA's asteroid explorer Hayabusa in late April this year.

As AKARI observed Itokawa on 26 July it was in the constellation of Scorpius, and was about 19 magnitudes bright in visible light. The asteroid and Earth were closest to each other, at a distance of about 42 million km (for comparison, Earth is 150 million km from the Sun). Given how close it was, Itokawa moved a significant distance on the sky over the short observing time.

Using observational data of asteroids such as Itokawa in combination with data from the explorer, models that estimate asteroid sizes can be made more accurate. This is especially useful for estimating the size of potentially hazardous asteroids which may be discovered in the future.

Before Hayabusa arrived at Itokawa, many observations to determine the asteroid's approximate size had already been attempted. Among the many different methods of measurement, the most accurate estimate was achieved by mid-infrared observations.

## Motion of Itokawa

With AKARI, it was possible to observe Itokawa at several different wavelengths in the mid-infrared range, obtaining a much more comprehensive set of data. This data is very important, not only for the study of the asteroid's infrared properties, but also for use as a template and source of comparison with other asteroids, to improve the estimates of their sizes.

Most sunlight falling on Itokawa is absorbed, heating the asteroid up. It then re-emits this energy as bright infrared light, which was in turn observed by AKARI. Only a small fraction of the incident sunlight is reflected from Itokawa, making it a very faint object when observed in visible light. It is very hard to observe using telescopes of sizes similar to that of AKARI from ground.

## **Positions of Itokawa and Earth**

Asteroid size is one of the most sought-after pieces of information. For asteroids that are not explored directly, their sizes can be estimated based on various observations from Earth. The temperature of asteroids is determined by the balance between the energy input from incident sunlight, and the output, emitted as infrared radiation.

Existing computer models estimate the temperature distribution in asteroids by considering their shape, rotational motion, and surface conditions.

Observational data in the mid-infrared gives information on the infrared light emitted by the asteroid. Asteroid size can be derived by comparing observational data in the mid-infrared, with that expected from the calculations of the model. The models can further be improved by using the infrared observational data of well-studied asteroids, such as Itokawa.

AKARI has also made observations of possible candidates for future asteroid exploration. It is expected that this detailed information will help greatly further our knowledge of these interesting relics of our Solar System.

Source: European Space Agency

Citation: AKARI's observations of asteroid Itokawa (2007, August 23) retrieved 25 April 2024 from <https://phys.org/news/2007-08-akari-asteroid-itokawa.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--