

Mice use specialized neurons to detect carbon dioxide in the air

August 16 2007

For mice, carbon dioxide often means danger - too many animals breathing in too small a space or a hungry predator exhaling nearby. Mice have a way of detecting carbon dioxide, and new research from Rockefeller University shows that a special set of olfactory neurons is involved, a finding that may have implications for how predicted increases in atmospheric carbon dioxide may affect animal behavior. The finding is reported in the August 17 issue of the journal *Science*.

Most olfactory sensory neurons express odorant receptor molecules and reside within the lining of the nasal cavity that detect odors. But a small subset express an enzyme called guanylyl cyclase-D (GC-D). Peter Mombaerts, professor and head of the Laboratory of Developmental Biology and Neurogenetics at Rockefeller, and Andreas Walz, a research associate in Mombaerts' lab, created a strain of mice in which GC-D expressing neurons glow with a green fluorescent protein. These GC-D expressing neurons also project their nerve endings to an unusual structure in the back of the olfactory bulb called necklace glomeruli, which resemble a string of beads.

The Rockefeller team's collaborators in China, led by Minmin Luo at the National Institute of Biological Sciences in Beijing, found that all the GC-D expressing neurons in the olfactory epithelium were activated by exposure to carbon dioxide. Conversely, all the cells in the lining of the nasal cavity that were activated by carbon dioxide are the GC-expressing neurons.

"These findings show that a specialized subsystem has evolved in the mouse to detect carbon dioxide," says Mombaerts.

Carbon dioxide makes up about four-hundredths of one percent of the atmosphere. To determine the threshold for carbon dioxide detection in the mouse, the Beijing team trained mice to lick water when they received a whiff of air with higher levels of carbon dioxide. As the amount of carbon dioxide in the whiffs of air was decreased, the accuracy of the animals' response became random. Statistical analysis fixed the threshold for detection at about six-hundredths of a percent, just above the average atmospheric level.

Mombaerts cautions that scientists still do not know if the GC-D enzyme is responsible for detecting carbon dioxide. "GC-D is a marker for the neurons that have specific carbon dioxide sensitivity," Mombaerts says. "At this time, we have not shown that the marker is mechanistically involved in sensory perception of carbon dioxide." The research does, however, suggest that scientists need to be on the lookout for behavioral changes in animals as carbon dioxide levels increase in the atmosphere, since, at some point increased levels may be detectable by animals, Mombaerts says.

"Animals may adapt to this gradual and persistent increase. Alternatively, the change may induce behavioral changes, such as an increase in irritability and aggression or a decrease in fertility," he says.

Source: Rockefeller University

Citation: Mice use specialized neurons to detect carbon dioxide in the air (2007, August 16)
retrieved 18 April 2024 from
<https://phys.org/news/2007-08-mice-specialized-neurons-carbon-dioxide.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.