

New, more direct pathways from outside the cell-to-cell nuclei discovered

August 10 2007

A team of Brooklyn College researchers has shattered a long-held belief that no direct pathway exists between material outside of a cell and the cell nucleus. (The cell is the smallest metabolically functional unit of life.)

It was already known that material outside of a cell can migrate into a cell. This occurs through processes known as endocytosis and phagocytosis, in which extracellular material is captured by a pinched-in segment of the cell membrane. This extracellular material then becomes trapped inside the resulting membrane-bound intracellular compartment, which is known as an endosome or phagosome.

It was also already known that material can migrate out of an endosome or phagosome and eventually enter the cell nucleus. But the Brooklyn College team has discovered that a phagosome and its contents can enter the cell's nucleus, where genetic information is stored and processed.

The team's discovery of the existence of direct pathways between extracellular material and cell nuclei will be published on August 10, 2007 in *Cell Motility and the Cytoskeleton*. The team was led by Ray Gavin and funded by the National Science Foundation (NSF).

Gavin says that these newly discovered direct pathways mean that "internalized material does not necessarily have to exit the phagosome before entering the nucleus." Therefore, his discovery means that "one less step is needed for extracellular material to get into the nucleus, and



so it is far easier than previously thought for this material to get into the nucleus."

Eve Barak, an NSF program director, describes Gavin's discovery as "an amazing and potentially paradigm-changing observation," and predicts that it "will have an enormous impact on how scientists think about how cellular functions are regulated."

Gavin has only observed direct pathways from the extracellular environment to the nucleus in the one-celled protozoan Tetrahymena thermophila. But Gavin says that "other discoveries that were based on this protozoan were later shown to be almost global in their occurrence in the biological world." Therefore, these direct pathways may operate in other organisms as well.

The discovery of the pathways in the protozoan resulted from an observation made by Gavin about 10 years ago while simply watching cells through a microscope. "I am a patient and constant observer of living things," he says. "And I watch them all the time with no agenda in mind. Some people are window gazers. They stand in front of the window and just look. I do the same with cells."

During one of Gavin's agenda-less cell-watching sessions, he noticed phagosomes clustered around a cell's nucleus. "It made me wonder why the phagosomes were positioned in that way," Gavin said. Thereafter, Gavin periodically looked for this intriguing behavior again, and occasionally glimpsed it again.

But Gavin remained unable to systematically search for phagosome clusters around nuclei until he received NSF funding in 2006 to purchase a confocal microscope, which provides three dimensional views of the cell. Just like a person would eventually find a worm hidden in an apple by repeatedly slicing through the apple, Gavin found the intruiging



phagosome behavior he sought by using the confocal microscope to repeatedly obtain views that sliced clear through the cell.

Specifically, the team tracked phagosomes carry extracellular material into the cell nuceli by introducing fluorescent latex beads into the area outside of the cell. They then observed the cells phagocytose (eat) the beads, which gradually moved to the nucleus. The arrival of the phagosomes and their loads at the nucleus was marked by the illumination of the nucleus by the beads. Similarly, the researchers also labeled the cellular membrane with a fluorescent dye, and then observed the pinched off, internalized membrane move to the nucleus.

"Biologists may now study the kinds of external molecules that can gain entry to the nucleus through these newly defined pathways and how these materials influence the nucleic material and its processes," said Gavin.

Source: National Science Foundation

Citation: New, more direct pathways from outside the cell-to-cell nuclei discovered (2007, August 10) retrieved 27 April 2024 from <u>https://phys.org/news/2007-08-pathways-cell-to-cell-nuclei.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.