

Truly sick or simply scared?

August 19 2007



Researchers at PNNL are developing a portable biomonitor to rapidly evaluate tiny samples of blood or saliva for exposure to nerve agents. Credit: Pacific Northwest National Laboratory

Scientists at Pacific Northwest National Laboratory have discovered a way to increase the sensitivity of test strips that will enable creation of a portable biosensor that can address a major concern associated with incidents involving chemical or nerve agents – the need to quickly distinguish between individuals who have been exposed and the “worried well.”

The sensor components resemble a pregnancy test strip and a small glucose testing meter. Its development will be discussed by principal investigator Yuehe Lin at the national meeting of the American

Chemical Society.

Every disease has biomarkers, a change in the proteins that announces something is wrong. Lin and his team are creating a nanoparticle “label” that can increase the ability of a sensor to detect and interpret the message of biomarkers.

“Current test strip based-immunoassay technology has very good selectivity, but it can only give a positive or negative response,” Lin said.

The researchers are working with an “electrochemical immunoassay approach.” This involves using the antibody of a specific disease — a protein produced in response to an invading bacterium or other foreign substance — to attract the biomarker. Lin found that labeling a second antibody with a nanoparticle amplifies the biomarker’s signal. Greater amplification means more precise readings.

Researchers at PNNL are developing a portable biomonitor to rapidly evaluate tiny samples of blood or saliva for exposure to nerve agents.

Lin achieves this by removing the iron from a nanoparticle-sized ball of the protein ferritin, creating an empty “cage” called apoferritin, which he then loads with another metal, such as cadmium. The cadmium-filled cage is attached to one end of the reporting antibody, and the immuno-reaction product becomes electroactive.

“The electrochemical signal is amplified several hundreds to thousand times because of the metal ions,” Lin said. “This level of sensitivity will allow detectors to be very precise in identifying the concentration of biomarkers in biological samples.”

Source: Pacific Northwest National Laboratory

Citation: Truly sick or simply scared? (2007, August 19) retrieved 20 April 2024 from <https://phys.org/news/2007-08-sick-simply.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.