


NASA finds Greenland snow melting hit record high in high places

September 25 2007




Microwave data from the Special Sensor Microwave Imaging radiometer was used to create this image of the 2007 Greenland melting anomaly which reflects the difference between the number of melting days occurring in 2007 and the average number of melting days during the period 1988 – 2006. Credit: NASA/Earth Observatory

A new NASA-supported study reports that 2007 marked an overall rise in the melting trend over the entire Greenland ice sheet and, remarkably,

melting in high-altitude areas was greater than ever at 150 percent more than average. In fact, the amount of snow that has melted this year over Greenland could cover the surface size of the U.S. more than twice.

Marco Tedesco, a research scientist at the Joint Center for Earth Systems Technology, cooperatively managed by NASA's Goddard Space Flight Center in Greenbelt, Md., and the University of Maryland at Baltimore County, used satellite data to compare average snow melting from 1988-2006 with what has taken place this summer.



Marco Tedesco summarizes the trend of the melting index in Greenland between 1988 and 2007 by showing a plot of how many surface size of the U.S. match the melting index observed each year of his study. Credit: NASA/Marco Tedesco and Luisa Valle

He found that in high altitude areas over 1.2 miles above sea level, the melting index -- an indicator of where melting is occurring and for how long - was significantly higher than average. Melting over those areas occurred 25-30 days longer this year than the observed average in the previous 19 years.

"When snow melts at those high altitudes and then refreezes, it can absorb up to four times more energy than fresh, unthawed snow," said Tedesco. "This can affect Earth's energy budget by changing how much radiation from the sun is absorbed by the Earth versus that reflected back into the atmosphere. Refrozen snow can also alter the snow density, thickness and snow-water content." Tedesco's findings were published Sept. 25 in the American Geophysical Union's Eos newspaper.

Researchers determine the melting index by multiplying how long melting took place by the area where the increased melting took place. According to Tedesco, melting in April and May of this year in high-altitude areas was very low, but in June melting jumped unexpectedly and led to the record melting index for the year.

"This record melting index in those areas came as a bit of a surprise, showing us, once again, the extreme variability and complexity of these processes," said Tedesco. His expertise in documenting melting trends produced other recent studies on increased snow melting over Greenland and the Antarctic.

The data collected by the Special Sensor Microwave Imagers on the Defense Meteorological Satellite Program satellites provided Tedesco with insight into how much of an electromagnetic signal was naturally emitted by snow and ice in areas beneath the satellite overpass. The microwave instruments can detect melting above and below the snow surface. The data were processed at the National Snow and Ice Data Center in Boulder, Colo., in just 24 hours after the satellite overpass,

enabling Tedesco to quickly spot changes that could signal a melting trend or new record.

Tedesco's work also confirmed that the melting index this year in lower altitude areas of Greenland, though not record breaking, was higher than average by 30 percent, placing 2007 in fifth place for the highest melting index after 2005, 2002, 1998 and 2004, in that order.

"Increases in the overall melting trend over Greenland have an impact that stretches beyond its icy shores," said Tedesco. "Aside from contributing to direct sea level rise, melting especially along the coast can speed up glaciers since the meltwater acts like a lubricant between the frozen surface and the bedrock deep below. The faster glaciers flow, the more water enters the ocean and potentially impacts sea level rise."

Tedesco, along with U.S. and European colleagues, plans to combine satellite data with related climate model results. He plans to visit Greenland in 2008 to compare his findings from space-borne data with those obtained by ground-based sensors, all with an eye to gathering further clues of what is happening on Greenland.

Source: Goddard Space Flight Center

Citation: NASA finds Greenland snow melting hit record high in high places (2007, September 25) retrieved 18 April 2024 from <https://phys.org/news/2007-09-nasa-greenland-high.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.