


Scientists Create Quantum Cascade Laser Nanoantenna

October 23 2007



The device consists of an optical antenna fabricated on the facet of a quantum cascade laser emitting infrared light with a wavelength of 7 microns. The Harvard team used nanofabrication techniques to form the optical antenna, which consists of two gold rectangles, each 1.2 microns long, separated by a narrow gap (100 nm). Light from the laser illuminates the antenna, resulting in an intense spot of light in the gap of size seventy times smaller than the wavelength. This is far smaller than what would be possible with the conventional approach of forming a spot of light by focusing with a lens. Due to the wave nature of light, such a spot would have a diameter of more than 7 microns. The figure is an electron microscope micrograph of the facet of the QC laser with the built-in nanoantenna. Shown are also an atomic force microscope topographic image of the antenna and an optical image obtained with a near field scanning optical microscope, showing the highly localized light spot in the antenna gap. Credit: Harvard University

In a major feat of nanotechnology engineering researchers from Harvard University have demonstrated a laser with a wide-range of potential applications in chemistry, biology and medicine. Called a quantum cascade (QC) laser nanoantenna, the device is capable of resolving the chemical composition of samples, such as the interior of a cell, with unprecedented detail.

Spearheaded by graduate students Nanfang Yu, Ertugrul Cubukcu, and Federico Capasso, Robert L. Wallace Professor of Applied Physics, all of Harvard's School of Engineering and Applied Sciences, the findings will be published as a cover feature of the October 22 issue of *Applied Physics Letters*. The researchers have also filed for U.S. patents covering this new class of photonic devices.

The laser's design consists of two gold rods separated by a nanometer gap (a device known as an optical antenna) built on the facet of a quantum cascade laser, which emits invisible light in the region of the spectrum where most molecules have their tell tale absorption fingerprints. The nanoantenna creates a light spot of nanometric size about fifty to hundred times smaller than the laser wavelength; the spot can be scanned across a specimen to provide chemical images of the surface with superior spatial resolution.

"There's currently a major push to develop powerful tabletop microscopes with spatial resolution much smaller than the wavelength that can provide images of materials, and in particular biological specimens, with chemical information on a nanometric scale," says Federico Capasso.

While infrared microscopes, based on the detection of molecular absorption fingerprints, are commercially available and widely used to map the chemical composition of materials, their spatial resolution is limited by the range of available light sources and optics to well above

the wavelength. Likewise the so-called near field infrared microscopes, which rely on an ultra sharp metallic tip scanned across the sample surface at nanometric distances, can provide ultrahigh spatial resolution but applications are so far strongly limited by the use of bulky lasers with very limited tunability and wavelength coverage.

"By combining Quantum Cascade Lasers with optical antenna nanotechnology we have created for the first time an extremely compact device that will enable the realization of new ultrahigh spatial resolution microscopes for chemical imaging on a nanometric scale of a wide range of materials and biological specimens," says Capasso.

Quantum cascade (QC) lasers were invented and first demonstrated by Capasso and his group at Bell Labs in 1994. These compact millimeter length semiconductor lasers, which are now commercially available, are made by stacking nanometer thick layers of semiconductor materials on top of each other. By varying the thickness of the layers one can select the wavelength of the QC laser across essentially the entire infrared spectrum where molecules absorb, thus custom designing it for a specific application.

In addition by suitable design the wavelength of a particular QCL can be made widely tunable. The range of applications of QC laser based chemical sensors is very broad, including pollution monitoring, chemical sensing, medical diagnostics such as breath analysis, and homeland security.

Source: Harvard University

Citation: Scientists Create Quantum Cascade Laser Nanoantenna (2007, October 23) retrieved 20 June 2024 from

<https://phys.org/news/2007-10-scientists-quantum-cascade-laser-nanoantenna.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.