


# A new view on sensing, movement, and behavioral control in animals

November 13 2007

---



A computer model of the knifefish illustrates the estimated SV for active sensing of prey (red) and stopping MV (blue). The backdrop shows a color map of the fish's simulated self-generated electric field. SV barely exceeds the stopping MV, revealing that the fish invests just enough energy into active sensing to detect prey in time to stop. Credit: MacIver et al.

While most animals, including humans, preferentially sense and move toward objects that are in front of them, an electric fish from the Amazon called the black ghost knifefish can swim backward or forward to catch its prey.

In a new study published online this week in the open-access journal *PLoS Biology*, James Snyder and colleagues at Northwestern University investigate the relationship between the energetic costs of the knifefish's active sensing system—which requires far more energy than passive sensing—and the area over which the animal senses its prey. They propose that the energetic constraints of the knifefish's active sensing system leads to a restricted sensory space compared to passive-sensing animals.

By combining video analysis of prey capture behavior with computational modeling of the fish's electrosensory capabilities, the scientists were able to quantify and compare the 3D volumes for sensation and movement for the first time in any animal.

They found that the sensory volume (the size and shape of the space within which objects can be detected by an animal) overlaps the motor volume (the location in space that an animal can reach within a set time period). They suggest that this coupling may arise from constraints that the animal faces when using self-generated energy to probe its environment. They also suggest that the degree of overlap between sensory and movement volumes can provide insight into the types of control strategies that are best suited for guiding behavior.

Citation: Snyder JB, Nelson ME, Burdick JW, MacIver MA (2007) Omnidirectional sensory and motor volumes in electric fish. *PLoS Biol* 5(11): e301. doi:10.1371/journal.pbio.0050301 ([www.plosbiology.org](http://www.plosbiology.org))

Source: Public Library of Science

Citation: A new view on sensing, movement, and behavioral control in animals (2007, November 13) retrieved 23 May 2024 from

<https://phys.org/news/2007-11-view-movement-behavioral-animals.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.