

# Recovering from a mass extinction

January 18 2008

---



The sabre-toothed Lycaenops was a top predator of the latest Permian in South Africa. Lycaenops was a gorgonopsian, one of a group of highly successful animals that dominated faunas in the Late Permian, but were wiped out, together with 90 per cent of all species, by the end-Permian mass extinction. Credit: Mike Benton, University of Bristol

**The full recovery of ecological systems, following the most devastating extinction event of all time, took at least 30 million years, according to new research from the University of Bristol**

The full recovery of ecological systems, following the most devastating extinction event of all time, took at least 30 million years, according to new research from the University of Bristol.

About 250 million years ago, at the end of the Permian, a major extinction event killed over 90 per cent of life on earth, including

insects, plants, marine animals, amphibians, and reptiles. Ecosystems were destroyed worldwide, communities were restructured and organisms were left struggling to recover. This was the nearest life ever came to being completely wiped out.

Previous work indicates that life bounced back quite quickly, but this was mostly in the form of ‘disaster taxa’ (opportunistic organisms that filled the empty ecospace left behind by the extinction), such as the hardy *Lystrosaurus*, a barrel-chested herbivorous animal, about the size of a pig.

The most recent research, conducted by Sarda Sahney and Professor Michael Benton at the University of Bristol and published in *Proceedings of the Royal Society B* this week, indicates that specialised animals forming complex ecosystems, with high biodiversity, complex food webs and a variety of niches, took much longer to recover.

Sahney said: “Our research shows that after a major ecological crisis, recovery takes a very long time. So although we have not yet witnessed anything like the level of the extinction that occurred at the end of the Permian, we should nevertheless bear in mind that ecosystems take a very long time to fully recover.”

Sahney and Benton looked at the recovery of tetrapods – animals with a backbone and four legs, such as amphibians and reptiles – and found that although globally tetrapods appeared to recover quickly, the dramatic restructuring that occurred at the community level was not permanent and communities did not recover numerically or ecologically until about 30 million years later.

Professor Benton explained: “Diversity is most commonly assessed by tallying the number of taxa on a global scale, but these studies are subject to the vagaries of sampling. By examining well-preserved and

well-studied faunas, the taxonomic and ecological recovery of communities after the Permian extinction event can be examined more accurately, and the problems of geological bias are largely avoided.”

The Permian extinctions occurred in three waves, the largest being at the boundary between the Permian and Triassic periods, 251 million years ago. This was the most devastating ecological event of all time, thought to be caused by large-scale volcanism in Russia which produced the ‘Siberian Traps’, covering over 200,000 square kilometers (77,000 square miles) in lava.

Source: University of Bristol

Citation: Recovering from a mass extinction (2008, January 18) retrieved 26 April 2024 from <https://phys.org/news/2008-01-recovering-mass-extinction.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.