

# Seeing through the dark

March 7 2008

---



A Dark Filament in Scattered Light  
(NTT/SOFI)

ESO Press Photo 06/08 (7 March 2008)

This image is copyright © ESO. It is released in connection with an ESO press release and may be used by the press on the condition that the source is clearly indicated in the caption.



Part of a filament in the Corona Australis molecular cloud. The image is a composite of J-, H-, and K-band near-infrared observations that were made with the SOFI instrument on ESO's NTT telescope in August 2006. The observations were made to test, how easily the scattered light can be observed and how good it is as a tracer of cloud structure. The J-, H-, and K-band intensities are coded with blue, green, and red colours. The gradual saturation of the near-infrared bands is visible as a change of colour. In diffuse regions the shorter wavelength J-band is strong and the colour is bluish. When the J-band saturates the colour changes first to green and finally, in the centre of the filament, the red colour corresponding to the K-band becomes the strongest. In the most saturated regions the surface brightness data can only be used to derive a lower limit for the total amount of dust on the line of sight. Credit: ESO

Astronomers have measured the distribution of mass inside a dark

filament in a molecular cloud with an amazing level of detail and to great depth. The measurement is based on a new method that looks at the scattered near-infrared light or 'cloudshine' and was made with ESO's New Technology Telescope. Associated with the forthcoming VISTA telescope, this new technique will allow astronomers to better understand the cradles of newborn stars.

The vast expanses between stars are permeated with giant complexes of cold gas and dust opaque to visible light. Yet these are the future nurseries of stars to be.

"One would like to have a detailed knowledge of the interiors of these dark clouds to better understand where and when new stars will appear," says Mika Juvela, lead author of the paper in which these results are reported.

Because the dust in these clouds blocks the visible light, the distribution of matter within interstellar clouds can be examined only indirectly. One method is based on measurements of the light from stars that are located behind the cloud.

"This method, albeit quite useful, is limited by the fact that the level of details one can obtain depends on the distribution of background stars," says co-author Paolo Padoan.

In 2006, astronomers Padoan, Juvela, and colleague Veli-Matti Pelkonen, proposed that maps of scattered light could be used as another tracer of the cloud's inner structure, a method that should yield more advantages. The idea is to estimate the amount of dust located along the line of sight by measuring the intensity of the scattered light.

Dark clouds are feebly illuminated by nearby stars. This light is scattered by the dust contained in the clouds, an effect dubbed 'cloudshine' by

Harvard astronomers Alyssa Goodman and Jonathan Foster. This effect is well known to sky lovers, as they create in visible light wonderful pieces of art called 'reflection nebulae'. The Chameleon I complex nebula is one beautiful example.

When making observations in the near-infrared, art becomes science. Near-infrared radiation can indeed propagate much farther into the cloud than visible light and the maps of scattered light can be used to measure the mass of the material inside the cloud.

To put this method to the test and use it for the first time for a quantitative estimation of the distribution of mass within a cloud, the astronomers who made the original suggestion, together with Kalevi Mattila, made observations in the near-infrared of a filament in the Corona Australis cloud. The observations were made in August 2006 with the SOFI instrument on ESO's New Technology Telescope at La Silla, in the Chilean Atacama Desert. The filament was observed for about 21 hours.

Their observations confirm that the scattering method is providing results that are as reliable as the use of background stars while providing much more detail.

"We can now obtain very high resolution images of dark clouds and so better study their internal structure and dynamics," says Juvela. "Not only is the level of details in the resulting map no longer dependent on the distribution of background stars, but we have also shown that where the density of the cloud becomes too high to be able to see any background stars, the new method can still be applied."

"The presented method and the confirmation of its feasibility will enable a wide range of studies into the interstellar medium and star formation within the Milky Way and even other galaxies," says co-author Mattila.

"This is an important result because, with current and planned near-infrared instruments, large cloud areas can be mapped with high resolution," adds Pelkonen. "For example, the VIRCAM instrument on ESO's soon-to-come VISTA telescope has a field of view hundreds of times larger than SOFI. Using our method, it will prove amazingly powerful for the study of stellar nurseries."

The report appears this week in the journal *Astronomy and Astrophysics* ("A Corona Australis cloud filament seen in NIR scattered light - I. Comparison with extinction of background stars", by Mika Juvela, Veli-Matti Pelkonen, Paolo Padoan, and Kalevi Mattila). Juvela, Pelkonen and Mattila are associated with the Helsinki University Observatory (Finland), while Padoan is at the University of California, San Diego, USA.

Source: ESO

Citation: Seeing through the dark (2008, March 7) retrieved 9 April 2024 from <https://phys.org/news/2008-03-dark.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--