

## **Researchers identify genes key to hormone production in plants**

April 3 2008



This scanning electron micrograph shows the abnormal anatomy of a group of developing flowers in an auxin-deficient wei8 tar2 double mutant plant. Credit: Jose Alonso, North Carolina State University

Researchers at North Carolina State University have pinpointed a small group of genes responsible for "telling" plants when, where and how to produce a hormone that is key to their development. Their findings shed light on the ways in which hormone production in plants affects both a plant's growth and its ability to adapt to changing environments.

Dr. Jose Alonso, assistant professor of genetics, and a team of geneticists



and plant biologists from NC State, Germany and the Czech Republic conducted the research. Their findings are published in the April 4 edition of the journal *Cell*.

Plant growth and development are regulated by a small number of hormones, which plants combine in various ways so that they can adapt to and thrive in changing environmental conditions. Auxin and ethylene are two of the most important of these growth-regulating hormones.

Scientists had previously established that plants respond differently to ethylene depending upon the type of plant tissue it is applied to, the developmental stage of the plant, and the surrounding environmental conditions. They also knew that the presence of auxin, another key growth-regulator, often served as a "trigger" for a plant to produce more ethylene, but were unsure of the ways in which auxin was synthesized.

"Auxin controls almost every process in a plant," Alonso says, "and so it's very important to understand how and why auxin is produced within the plant."

In order to find out more about how auxin production is triggered, the NC State team identified a mutant strain of Arabidopsis – or mustard weed – that had a root system insensitive to the growth inhibitory effect of ethylene.

When the team looked at the genome of this mutant strain of mustard weed, they discovered that its lack of response to ethylene was due to the changes in a gene that they named TAA1. This gene produces a protein that is necessary for auxin synthesis. In a normal plant, the TAA1 gene recognizes the presence of ethylene as its signal to make proteins that in turn synthesize auxin, which controls growth.

The researchers found that if the TAA1 gene and two other related genes



were "knocked out" or inactive, the plant had 50 percent less auxin than normal.

Their findings are the first to definitively establish a relationship between a particular family of genes, tissue-specific ethylene response, and auxin production in plants.

"If we want to do intelligent manipulation of plants, to breed them so that they ripen at a certain rate, or so that they're well-adapted to particular environments, then we need to understand more about the ways that these hormones interact or 'talk' to each other," Alonso says. "This research gives us concrete evidence for at least one way in which this happens."

Source: North Carolina State University

Citation: Researchers identify genes key to hormone production in plants (2008, April 3) retrieved 2 May 2024 from <u>https://phys.org/news/2008-04-genes-key-hormone-production.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.