

# Large methane release could cause abrupt climate change as happened 635 million years ago

May 28 2008

---



Dolomite cement, formed from oxidized methane as it evolved from melting methane hydrates at the end of the snowball Earth glaciation, present in wave-cut platforms at Marino Rocks, South Australia. The dolomite is orange-red and formed vertical plumbing of tubes and vugs as methane passed upward and disrupted overlying sediment. Credit: M. Kennedy, UC Riverside

An abrupt release of methane, a powerful greenhouse gas, about 635 million years ago from ice sheets that then extended to Earth's low latitudes caused a dramatic shift in climate, triggering a series of events that resulted in global warming and effectively ended the last "snowball" ice age, a UC Riverside-led study reports.

The researchers posit that the methane was released gradually at first and

then in abundance from clathrates – methane ice that forms and stabilizes beneath ice sheets under specific temperatures and pressures. When the ice sheets became unstable, they collapsed, releasing pressure on the clathrates which began to degas.

“Our findings document an abrupt and catastrophic means of global warming that abruptly led from a very cold, seemingly stable climate state to a very warm also stable climate state with no pause in between,” said Martin Kennedy, a professor of geology in the Department of Earth Sciences, who led the research team.

“This tells us about the mechanism, which exists, but is dormant today, as well as the rate of change,” he added. “What we now need to know is the sensitivity of the trigger: how much forcing does it take to move from one stable state to the other, and are we approaching something like that today with current carbon dioxide warming.”

Study results appear in the May 29 issue of *Nature*.

According to the study, methane clathrate destabilization acted as a runaway feedback to increased warming, and was the tipping point that ended the last snowball Earth. (The snowball Earth hypothesis posits that the Earth was covered from pole to pole in a thick sheet of ice for millions of years at a time.)

“Once methane was released at low latitudes from destabilization in front of ice sheets, warming caused other clathrates to destabilize because clathrates are held in a temperature-pressure balance of a few degrees,” Kennedy said. “But not all the Earth’s methane has been released as yet. These same methane clathrates are present today in the Arctic permafrost as well as below sea level at the continental margins of the ocean, and remain dormant until triggered by warming.

“This is a major concern because it’s possible that only a little warming can unleash this trapped methane. Unzipping the methane reservoir could potentially warm the Earth tens of degrees, and the mechanism could be geologically very rapid. Such a violent, zipper-like opening of the clathrates could have triggered a catastrophic climate and biogeochemical reorganization of the ocean and atmosphere around 635 million years ago.”

Today, the Earth’s permafrost extends from the poles to approximately 60 degrees latitude. But during the last snowball Earth, which lasted from 790 to 635 million years ago, conditions were cold enough to allow clathrates to extend all the way to the equator.

According to Kennedy, the abruptness of the glacial termination, changes in ancient ocean-chemistry, and unusual chemical deposits in the oceans that occurred during the snowball Earth ice age have been a curiosity and a challenge to climate scientists for many decades.

“The geologic deposits of this period are quite different from what we find in subsequent deglaciation,” he said. “Moreover, they immediately precede the first appearance of animals on earth, suggesting some kind of environmental link. Our methane hypothesis is capable also of accounting for this odd geological, geochemical and paleoceanographic record.”

Also called marsh gas, methane is a colorless, odorless gas. As a greenhouse gas, it is about 30 times more potent than carbon dioxide, and has largely been held responsible for a warming event that occurred about 55 million years ago, when average global temperatures rose by 4-8 degrees Celsius.

When released into the ocean-atmosphere system, methane reacts with oxygen to form carbon dioxide and can cause marine dysoxia, which

kills oxygen-using animals, and has been proposed as an explanation for major oceanic extinctions.

“One way to look at the present human influence on global warming is that we are conducting a global-scale experiment with Earth’s climate system,” Kennedy said. “We are witnessing an unprecedented rate of warming, with little or no knowledge of what instabilities lurk in the climate system and how they can influence life on Earth. But much the same experiment has already been conducted 635 million years ago, and the outcome is preserved in the geologic record. We see that strong forcing on the climate, not unlike the current carbon dioxide forcing, results in the activation of latent controls in the climate system that, once initiated, change the climate to a wholly different state.”

As part of their research, Kennedy and his colleagues collected hundreds of marine sediment samples in South Australia for stable isotope analysis, an important tool used in climate reconstruction. At UCR, the researchers analyzed the samples and found the broadest range of oxygen isotopic variation ever reported from marine sediments that they attribute to melting waters in ice sheets as well as destabilization of clathrates by glacial meltwater.

Next in their research, Kennedy and his colleagues will work on estimating how much of the temperature change that occurred 635 million years ago was due solely to methane.

Source: University of California - Riverside

Citation: Large methane release could cause abrupt climate change as happened 635 million years ago (2008, May 28) retrieved 5 April 2024 from <https://phys.org/news/2008-05-large-methane-abrupt-climate-million.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.