


# Using a light touch to measure protein bonds

June 30 2008

---



The strength of actin binding protein interactions cross-linking a surface bound and bead tethered actin filament are probed using force from an optical trap. Filamin in green which forms networks of actin filaments shown as a confocal image on the right and alpha-actinin in blue which bundles actin filaments, shown as an image on the left, were probed using this assay configuration. Image courtesy / Hyungsuk Lee, Jorge M. Ferrer, and Matthew J. Lang

MIT researchers have developed a novel technique to measure the strength of the bonds between two protein molecules important in cell machinery: Gently tugging them apart with light beams.

"It's really giving us a molecular-level picture of what's going on," said Matthew Lang, an assistant professor of biological and mechanical engineering and senior author of a paper on the work appearing in the June 30 advanced online issue of the *Proceedings of the National*

*Academy of Sciences.*

Last fall, Lang and others demonstrated that light beams could be used to pick up and move individual cells around the surface of a microchip.

Now they have applied the optical tweezers to measuring protein microarchitectures, allowing them to study the forces that give cells their structure and the ability to move.

The researchers focused on proteins that bind to actin filaments, an important component of the cytoskeleton. Depending on the arrangement and interaction of actin filaments, they can provide structural support, help the cell crawl across a surface or sustain a load (in muscle cells).

"We're trying to understand how nature engineered these molecular linkages to use in different ways," said Lang.

Actin filaments are most commonly found either bonded or crosslinked by a much smaller actin binding protein.

The researchers studied the interactions between the proteins by pinning one actin filament to a surface and controlling the motion of the second one with a beam of light. As the researchers tug on a bead attached to the second filament, the bond mediated by the actin-binding protein eventually breaks.

With this technique, the researchers can get a precise measurement of the force holding the proteins together, which is on the order of piconewtons ( $10^{-12}$  newtons).

The same technique could be used to investigate many of the other hundreds of protein interactions that occur in the cytoskeleton, said

Lang.

Source: Massachusetts Institute of Technology

Citation: Using a light touch to measure protein bonds (2008, June 30) retrieved 20 March 2024 from <https://phys.org/news/2008-06-protein-bonds.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.