

# Discovery of a mechanism that regulates cell movement

July 20 2008

---

A study performed by researchers at the Institute for Research in Biomedicine (IRB Barcelona), in collaboration with researchers at the Instituto de Biología Molecular of the CSIC, reveal a mechanism that controls the movement of cells in a tissue by regulating cell adhesion. This same mechanism may be defective in diseases such as cancer and its metastasis, when tumour cells lose their adhesion to neighbouring cells and migrate through the organism. The results of this research have been published in this week's *Nature Cell Biology*.

Specialists in development, Daniel Shaye, Jordi Casanova and Marta Llimargas, have studied the mechanisms that control cell movements during trachea development in the fly *Drosophila*. In this process, cells that initially form column of two cells deep, change their position to line up one after the other in a single file.

A key element in the regulation of these movements is the amount of adhesive protein E-Cadherin located in the cellular membrane. Jordi Casanova, head of the Morphogenesis in *Drosophila* at IRB Barcelona, explains that "when movement starts, the levels of this protein in the cells decrease, thereby allowing them to slide one on top of the other, and once in this position the levels of this protein are re-established in order to seal the new binding alignment".

The in-depth study of this phenomenon has led to the finding that the amount of E-Cadherin on the cell surface is controlled by the trafficking of this protein inside the cell and the identification of the elements that

regulate this transport. During the experiments, the researchers induced or blocked cell movement by modifying the elements that control the trafficking of adhesive protein towards the membrane.

"We demonstrate a mechanism that explains how cells can change their position within a given tissue. The sequence is clear: the greater the amount of protein, the greater the adhesion and the smaller the movement", recaps Casanova.

## **A mechanism may be dysfunctional during cancer progression and metastases**

The role of E-Cadherin in the binding between epithelial cells is universal in all animals and therefore, following logic, the researchers believe that the mechanisms that regulate the levels of this protein may also be universal.

"We speculate that the regulatory mechanism that we have discovered may also be present in other developmental contexts. However, in addition, fundamental elements of the mechanism may show a dysfunction in pathological processes such as the progression of cancer and metastases", adds Casanova.

As revealed by this same researcher, the amounts of E-Cadherin and of one of the elements required for trafficking decrease in the progression of a kind of oesophagus cancer and this effect is related to the gain in tumour cell motility, which allows them to spread throughout the body more rapidly.

Thanks to this study, researchers working on cancer will be able to use animal models to test whether this mechanism is defective in other kinds of tumours and to study whether any of its genetic components are valid

as therapeutic targets.

Source: Institute for Research in Biomedicine

Citation: Discovery of a mechanism that regulates cell movement (2008, July 20) retrieved 8 May 2024 from <https://phys.org/news/2008-07-discovery-mechanism-cell-movement.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.