

## Will our future brains be smaller?

July 9 2008



The speed at which we react to threatening situations can have life or death implications. In the more primitive past, it could have meant escaping a wild animal; today it might mean swerving to avoid a head-on car crash.

It has been thought for some years that mammals have two decisionmaking systems in their brains which operate at different speeds to cope with different situations. New research from the University of Bristol supports this theory and has shown that the evolutionary pressures arising from the older, faster, but less accurate, part of the brain may have shaped the more recent development of the slower-acting but more precise cortex, found in humans and higher animals. The research is published today in *Proceedings of the Royal Society B*.



Pete Trimmer, lead author on the study, explained: "If we compare the brain of a human with that of a reptile, we find they are very similar except that mammals have a large 'outer cortex' around the outside of the existing 'sub-cortical' brain, that is common to other vertebrates.

"The fact that lizards make decisions indicates that the sub-cortical brain in humans is also likely to be used in decision-making. However, fMRI scans now reveal that parts of the outer cortex (which developed more recently in our evolutionary past) are also used when making decisions."

Why does the brain need these two decision-making areas? What benefit does the new cortex bring? After all, extra brain means extra weight and energy required to carry it around. Furthermore, is the older sub-cortical system now largely redundant? If so, could we expect it to atrophy in future humans so our brains become smaller?

To address these questions, Trimmer built theoretical models representing the two systems in which the sub-cortical system was assumed to act very quickly but inaccurately, whereas the cortex allowed information to be gathered before making an informed decision, and was therefore slower.

The results of their modelling showed that when the threat level is high, such as the risk of being attacked by a dangerous animal, it is very useful to have the fast-acting, if inaccurate, system. But when dealing with situations which don't occur very often, or complex scenarios with many conflicting cues such as social situations, the cortical system is of more use than the sub-cortical system.

Trimmer commented: "As life became more complex, the benefit of gathering information before making a decision put an evolutionary pressure on the early brain. This may have led to the rapid development of the cortex in mammals. So if humans continue to live in a world of



dangers such as wild animals or fast-moving cars, there will still be an evolutionary benefit to maintaining the sub-cortical system, and it is unlikely to atrophy in future humans."

Source: University of Bristol

Citation: Will our future brains be smaller? (2008, July 9) retrieved 20 May 2024 from <u>https://phys.org/news/2008-07-future-brains-smaller.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.