

Large dead zones predicted for Gulf, Chesapeake Bay

July 14 2008

Record-setting "dead zones" in the Gulf of Mexico and Chesapeake Bay appear likely this summer, according to new forecasts from a University of Michigan researcher.

Donald Scavia, a professor at the U-M School of Natural Resources and Environment (SNRE), makes the annual forecasts using models driven by nutrient load estimates from the U.S. Geological Survey.

In this year's Chesapeake Bay Hypoxic Volume Forecast, Scavia predicts a summer hypoxic volume of 9.9 cubic kilometers (2.4 cubic miles), the sixth-highest on record. If the upper value of the forecast range of 12.3 cubic kilometers (3 cubic miles) is reached, it will be the highest on record. The Bay is about 200 miles long on the East Coast and stretches from Maryland to Virginia. It supports thousands of species of plants, fish and animals. The Bay's oxygen levels are critical in determining the health of its ecosystem.

Given recent massive flooding of cities and farms in the Mississippi River basin, the Gulf of Mexico Hypoxia Area Forecast is for the dead zone to cover between 21,500 and 22,500 square kilometers (8,400-8,800 square miles) of bottom waters along the Louisiana-Texas coast. If the prediction bears out, it will be the largest on record.

"The growth of these dead zones is an ecological time bomb," said Scavia, who is also director of the Michigan Sea Grant program based at SNRE. "Without determined local, regional, and national efforts to

control them, we are putting major fisheries at risk." According to Scavia, the best way to shrink the dead zones is to reduce the amount of nitrogen and phosphorous flowing into these water basins.

Hypoxia refers to the loss of oxygen in water, which then leads to conditions unsustainable for aquatic life. The Chesapeake Bay and Gulf of Mexico dead zones form each spring as nitrogen and phosphorus loads from farm fertilizers, atmospheric deposition and wastewater treatment plants stimulate algae blooms. These blooms eventually die and sink to the bottom, where bacteria decompose them and consume most of the oxygen. The zones dissipate each fall as changes in water currents and temperatures mix and reaerate the water—only to return the next spring.

Scavia originally developed the model to estimate the nitrogen-load reductions needed to reach particular hypoxia goals. Several years ago, he discovered that the model could also forecast the dead zone size for an upcoming season, based on the average January-May nitrogen loads for the Chesapeake Bay, and average May-June loads for the Mississippi River basin.

Source: University of Michigan

Citation: Large dead zones predicted for Gulf, Chesapeake Bay (2008, July 14) retrieved 9 April 2024 from <https://phys.org/news/2008-07-large-dead-zones-gulf-chesapeake.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
