

# DNA editing tool flips its target

September 3 2008

---

Imagine having to copy an entire book by hand without missing a comma. Our cells face a similar task every time they divide. They must duplicate both their DNA and a subtle pattern of punctuation-like modifications on the DNA known as methylation.

Scientists at Emory University School of Medicine have caught in action one of the tools mammalian cells use to maintain their pattern of methylation. Visualized by X-ray crystallography, the SRA domain of the protein UHRF1 appears to act like a bookmark while enzymes are copying a molecule of DNA.

The team's description of the protein's structure while bound to DNA is published this week in *Nature*.

Scientists refer to methylation, the addition of a methyl group to DNA, as an "epigenetic" modification because it adds a layer of information on top of the genetic sequence of the DNA itself. It marks genes for silencing, which means they do not manufacture proteins.

"The processes that copy the methylation pattern have to be faithful," says senior author Xiaodong Cheng, PhD, professor of biochemistry and a Georgia Research Alliance eminent scholar. "Otherwise, losing DNA methylation marks can have serious consequences, causing genes to become active at the wrong places and times."

"Gene silencing via DNA methylation is critical for normal development and for curbing the runaway cell division that characterizes cancer," said

Peter Preusch, PhD, who oversees biophysics grants at the National Institute of General Medical Sciences of the National Institutes of Health. "Alterations in methylation patterns are also important for generating embryonic stem-like cells from differentiated cells."

In mammalian cells, methylation usually appears on double stranded DNA where the nucleotide Cytosine (C) is followed by Guanine (G). The complementary sequence on the opposite strand is also C then G, and the methylation appears on both Cs.

When a cell is copying its DNA, a set of enzymes duplicates the DNA sequence from the parental strand to the new "daughter" strand but not the methylation. Each new daughter strand of the DNA molecule is left with the previously methylated Cs unmethylated. UHRF1 recognizes this "hemi-methylated" DNA and calls in a methyltransferase enzyme to add a second methyl group onto the daughter strand.

"UHRF1 has the important task of making sure the methyltransferase enzyme does its job in the right place and right time," Cheng says.

Mouse cells that have deleted the UHRF1 gene are more sensitive to DNA-damaging agents such as radiation, and mouse embryos without the gene cannot complete development. Other studies have found that cancer cells produce more UHRF1 than non-cancerous cells.

What was an unexpected finding was how the SRA domain of UHRF1 recognizes the hemi-methylated DNA, Cheng says. It flips the methylated nucleotide out of the DNA helix, which only had been seen previously in enzymes that physically modify the DNA.

Cheng says the flipping mechanism could prevent the protein from sliding away once it has found a hemi-methylated site.

"It suggests that it serves as a placeholder, where it recruits other enzymes for faithful DNA methylation or repair enzymes if the DNA has been damaged," he says.

Source: Emory University

Citation: DNA editing tool flips its target (2008, September 3) retrieved 27 April 2024 from <https://phys.org/news/2008-09-dna-tool-flips.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.