

Just a little squeeze lets proteins assess DNA

December 16 2008



Photograph of crystals of the DNA-snipping protein HincII taken under a microscope. The crystals are iridescent because they refract light. The longest crystal is about one millimeter. Photo credit: Nancy C. Horton. Credit: Nancy C. Horton, The University of Arizona

(PhysOrg.com) -- To find its target, all a protein needs to do is give quick squeezes as it moves along the DNA strand, suggests new research from The University of Arizona in Tucson.

Scientists had thought DNA-binding proteins primarily used full-body hugs for accurate readings of the information coded in the DNA's sequence.

Even a protein known to use the hug method, called direct readout, can

effectively pinpoint sites on DNA using indirect readout, found researcher Nancy C. Horton and her colleagues.

"It was a total surprise," said Horton, a UA associate professor of biochemistry and molecular biophysics. "No one had ever seen it before."

Doing the quick squeezes that scientists call indirect readout probably works faster than requiring full-body contact with all the DNA, the researchers suggest. Quick and accurate identification of key sites on DNA is important for the health of all kinds of cells, from bacteria to humans.

To detect the protein-DNA connection in such detail, Horton and her co-authors Elizabeth J. Little and Andrea C. Babic studied a DNA-binding protein that bacteria use to protect themselves from viral infections.

The finding has implications for the development of designer drugs.

"People have and are developing DNA-binding proteins to turn genes on and off," Horton said. Such designer proteins can be used to cut out the bad copy of a gene and help replace it with good copy.

"We found that indirect readout is important for finding the right sequence, and we now think indirect readout is also important for finding it quickly," she said.

The team published their paper, "Early Interrogation and Recognition of DNA Sequence by Indirect Readout," in the December issue of the journal *Structure*. First author Little and co-author Babic were postdoctoral research associates in Horton's laboratory when they did the research. The two are now senior scientists at Ventana Medical Systems, Inc. in Tucson, Ariz.

The National Institutes of Health funded the research.

Horton studies proteins that bind to DNA.

Seven years ago, she figured out the structure of a protein called HincII that snips up DNA. The protein is a type of enzyme called a restriction endonuclease and comes from *Haemophilus influenzae* bacteria.

Since that time, Horton has been trying to learn how HincII interrogates the DNA to find the right place to cut.

The protein protects bacteria by cutting up DNA from invading viruses. Without the protective protein, viral DNA would commandeer the bacterium's cellular machinery to produce viruses and ultimately kill the bacterial cell.

The HincII protein distinguishes between bacterial DNA and viral DNA by recognizing certain sequences on DNA. Such a defense requires speed to prevent the marauding virus from killing the cell and also accuracy so the protein doesn't accidentally hack up the bacterium's own DNA.

Horton knew from her previous work that the HincII protein used the direct readout method to find the particular sequence of DNA that corresponded to enemy DNA. The protein seemed to distort the DNA to read it.

Removing the direct readout contact between the protein and the DNA might show whether the DNA distortion or the contact itself was important, Horton said.

Therefore Little and Babic created a mutant protein that couldn't hug DNA closely and therefore couldn't use the direct readout method. Little

described the mutant protein as missing the fingers the normal protein used to probe the DNA.

"If the finger was doing all the recognition, then the mutant should cut any DNA sequence," Horton said.

To see how the mutant interacted with DNA, the researchers crystallized the mutant protein-DNA complex in action.

Initially, Horton thought the assay had gone wrong and almost threw the results in the trash, she wrote in an e-mail.

The mutant protein had chosen the proper site on the DNA with 100 percent specificity, which was opposite from what she expected. In addition, the DNA was distorted, even though the mutant couldn't make the strong contact a normal protein would.

"I did a double-take. I was just taking a picture to have a record that it was non-specific," she said.

Understanding how endonucleases and other DNA binding-proteins recognize a particular DNA sequence provides insight into key cellular processes including the replication, transcription and repair of DNA.

Little said, "In every single one of your cells are proteins looking for the proper sequences in DNA in order to make the proteins you need to stay alive."

Horton added, "Understanding how these processes work helps in the understanding of diseases so that we could potentially cure the disease."

Source: University of Arizona

Citation: Just a little squeeze lets proteins assess DNA (2008, December 16) retrieved 20 April 2024 from <https://phys.org/news/2008-12-proteins-dna.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.