

# Nanoparticles Double Their Chances of Getting Into Sticky Situations

February 16 2009

---



Stefan Bon and David Cheung with an image from their paper

(PhysOrg.com) -- Chemistry researchers at the University of Warwick have found that tiny nanoparticles could be twice as likely to stick to the interface of two non mixing liquids than previously believed. This opens up a range of new possibilities for the uses of nanoparticles in living cells, polymer composites, and high-tech foams, gels, and paints. The researchers are also working on ways of further artificially enhancing this new found sticking power.

In a paper entitled "Interaction of nanoparticles with ideal liquid-liquid interfaces" just published in *Physical Review Letters* the University of Warwick researchers reviewed molecular simulations of the interaction between a non-charged nanoparticle and an "ideal" liquid-liquid interface. They were surprised to find that very small nanoparticles (of around 1 to 2 nanometres) varied considerably in their simulated ability

to stick to such interfaces from what was expected in the standard model.

The researchers found that it took up to 50 percent more energy to dislodge the particles from the liquid-liquid interface for the smallest particle sizes. However as the radius of the particles increased this deviation from the standard model gradually faded out.

The researchers, Dr ir Stefan A. F. Bon and Dr David L. Cheung, believe that previous models failed to take into account the action of "capillary waves" in their depiction of the nanoparticles behaviour at the liquid to liquid interfaces.

Dr ir Stefan A. F. Bon said

"This new understanding on the nano-scale gives us much more flexibility in the design of everything from high-tech composite materials, to the use of quantum dots, cell biochemistry, and the manufacture of new "armored" polymer paint particles."

The researchers are now working on ways to build on this newly found natural stickiness of nanoparticles by designing polymer nanoparticles with opposing hydrophobic and hydrophilic surfaces that will bind even more strongly at oil/water liquid interfaces.

Provided by University of Warwick

Citation: Nanoparticles Double Their Chances of Getting Into Sticky Situations (2009, February 16) retrieved 18 April 2024 from <https://phys.org/news/2009-02-nanoparticles-chances-sticky-situations.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.