

Colors of Quasars Reveal a Dusty Universe

February 26 2009

(PhysOrg.com) -- The vast expanses of intergalactic space appear to be filled with a haze of tiny, smoke-like "dust" particles that dim the light from distant objects and subtly change their colors, according to a team of astronomers from the Sloan Digital Sky Survey (SDSS-II), including a researcher from the University of California, Davis.

SDSS-II is the second step of the Sloan Digital Sky Survey, which since 2000 has used a 2.5-meter telescope coupled with a 120-megapixel camera and two spectrographs to probe the night sky from an observatory in New Mexico.

"Galaxies contain lots of dust, most of it formed in the outer regions of dying stars," said team leader Brice Ménard of the Canadian Institute for Theoretical Astrophysics. "The surprise is that we are seeing dust hundreds of thousands of light-years outside of the galaxies, in intergalactic space."

The new findings are reported in a paper titled "Measuring the galaxy-mass and galaxy-dust correlations through magnification and reddening," submitted to the journal *Monthly Notices of the Royal Astronomical Society*, and posted today at arxiv.org/abs/0902.4240 .

To discover this intergalactic dust, the team analyzed the colors of distant quasars whose light passes in the vicinity of foreground galaxies on its way to the Earth.

Dust grains block blue light more effectively than red light, explained

astronomer Ryan Scranton of UC Davis, another member of the discovery team. "We see this when the sun sets: light rays pass through a thicker layer of the atmosphere, absorbing more and more blue light, causing the sun to appear reddened."

"We find similar reddening of quasars from intergalactic dust, and this reddening extends up to 10 times beyond the apparent edges of the galaxies themselves," said Scranton, who is an assistant researcher in the physics department.

The team analyzed the colors of about 100,000 distant quasars located behind 20 million galaxies, using images from SDSS-II. "Putting together and analyzing this huge data set required cutting-edge ideas from computer science and statistics," said team member Gordon Richards of Drexel University. "Averaging over so many objects allowed us to measure an effect that is much too small to see in any individual quasar."

Supernova explosions and "winds" from massive stars drive gas out of some galaxies, Ménard explained, and this gas may carry dust with it. Alternatively, the dust may be pushed directly by starlight.

"Our findings now provide a reference point for theoretical studies," said Ménard.

Intergalactic dust could also affect planned cosmological experiments that use supernovae to investigate the nature of "dark energy," a mysterious cosmic component responsible for the acceleration of the expansion of the universe.

"Just like household dust, cosmic dust can be a nuisance," said Scranton. "Our results imply that most distant supernovae are seen through a bit of haze, which may affect estimates of their distances."

Intergalactic dust doesn't remove the need for dark energy to explain current supernova data, Ménard explained, but it may complicate the interpretation of high-precision distance measurements. "These experiments are very ambitious in their goals," said Ménard, "and subtle effects matter."

Provided by University of California, Davis

Citation: Colors of Quasars Reveal a Dusty Universe (2009, February 26) retrieved 16 June 2024 from <https://phys.org/news/2009-02-quasars-reveal-dusty-universe.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--