

Birds in Flint Hills of Kansas, Oklahoma face population decline despite large habitat

March 3 2009



A dickcissel, eastern meadowlark and grasshopper sparrow are three bird species common to the Flint Hills of Kansas and Oklahoma. A Kansas State University ecologist and her colleagues found these birds experiencing serious population decline in the face of extensive land-management practices like annual burning and widespread grazing. Credit: David A. Rintoul, K-State Division of Biology

The wide-open spaces of the Flint Hills may no longer provide a secure home on the range for several familiar grassland birds, according to research by a Kansas State University ecologist and her colleagues.

The researchers found that three bird species common to the Flint Hills region of Kansas and Oklahoma are experiencing serious population decline in the face of extensive land-management practices like annual burning and widespread grazing

The Flint Hills contain the largest remaining tracts of tallgrass prairie, a habitat that once covered much of the Great Plains but of which only 4 percent is now left, mostly in the Flint Hills. Far from being pristine

prairie, however, the Flint Hills region supports a major cattle industry and is intensively managed.

"Because of its size, the Flint Hills is assumed to be a population stronghold for grassland birds," said Kimberly With, a K-State associate professor of biology who led the study. "Mostly this has been based on bird counts, but they can be misleading because they don't show what the region is capable of producing. Birds are very mobile and thus birds could come from elsewhere to give the appearance of a stable population year after year. This is especially true if the region attracts birds because of its size, but birds do not breed successfully once they settle here."

With was joined by ecologists Anthony King from the Oak Ridge National Laboratory and William Jensen, a former postdoctoral researcher in With's lab at K-State. They conducted a two-year study of regional viability of three grassland birds: the dickcissel, grasshopper sparrow and eastern meadowlark. With and her colleagues found that none of these bird species is viable in the 4 million-acre Flint Hills region. They estimated population declines of as much as 29 percent per year during the years studied.

The results were published in December 2008 in the journal *Biological Conservation*, <http://tinyurl.com/cza3xy>

"Although considered to be relatively common, all three of these species have been exhibiting declines at a continent-wide level," With said.

According to a 2007 report by The National Audubon Society, two of these species -- the grasshopper sparrow and eastern meadowlark -- have lost 62 percent and 75 percent, respectively, of their global population in the past 40 years.

With and her colleagues found that birds are not breeding successfully in

the Flint Hills. With said that more than 80 percent of nests were destroyed by predators. They also found that birds nesting in native prairie hayfields were more successful than birds nesting in grazed grassland. She said this might be because in the Flint Hills, the hayfields were mown later than in other prairie regions, enabling birds to finish nesting before the nests are mowed under.

"These hayfields are a less disturbed habitat than other managed grasslands in the Flint Hills, 90 percent of which is grazed and up to two-thirds may be burned," With said.

The researchers suggest that land-management practices may offer only part of the explanation for declining bird populations.

"There was a drought one of the two years we studied the birds, and in that year birds didn't do well anywhere," With said. "Under global climate change, we may see more prolonged droughts and less frequent, but more intense storms. If rain comes as intense storms with hail, that could be just as lethal to the birds."

With said that more study is needed before recommendations can be made to help manage these bird populations. But she said that the results are still valuable because -- like a canary in a coal mine -- birds indicate a region's overall environmental health.

"The cattle industry is important economically to the region," With said. "The region is also ecologically important. The challenge is to what extent you can successfully achieve these joint aims of agriculture and conservation within the Flint Hills."

Source: Kansas State University

Citation: Birds in Flint Hills of Kansas, Oklahoma face population decline despite large habitat (2009, March 3) retrieved 1 May 2024 from <https://phys.org/news/2009-03-birds-flint-hills-kansas-oklahoma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.