

## New experiments constrain Higgs mass (w/Videos)

## March 13 2009



Scientists from the CDF and DZero collaborations at DOE's Fermilab have combined Tevatron data from their two experiments to increase the sensitivity for their search for the Higgs boson. While no Higgs boson has been found yet, the results announced today exclude a mass for the Higgs between 160 and 170  $GeV/c^2$  with 95 percent probability. A larger area is excluded at the 90 percent probability level. Earlier experiments at the Large Electron-Positron Collider at CERN excluded a Higgs boson with a mass of less than 114 GeV/c<sup>2</sup> at 95 percent probability. Calculations of quantum effects involving the Higgs boson require its mass to be less than 185 GeV/c<sup>2</sup>. The results show that CDF and DZero are sensitive to potential Higgs signals. The Fermilab experimenters will test more and more of the available mass range for the Higgs as their experiments record more collision data and as they continue to refine their experimental analyses.

(PhysOrg.com) -- The territory where the Higgs boson may be found continues to shrink. The latest analysis of data from the CDF and DZero collider experiments at the U.S. Department of Energy's Fermilab now excludes a significant fraction of the allowed Higgs mass range



established by earlier measurements. Those experiments predict that the Higgs particle should have a mass between 114 and 185  $\text{GeV/c}^2$ . Now the CDF and DZero results carve out a section in the middle of this range and establish that it cannot have a mass in between 160 and 170  $\text{GeV/c}^2$ .

"The outstanding performance of the <u>Tevatron</u> and CDF and DZero together have produced this important result," said Dennis Kovar, Associate Director of the Office of Science for <u>High Energy Physics</u> at the U.S. Department of Energy. "We're looking forward to further Tevatron constraints on the Higgs <u>mass</u>."

The Higgs particle is a keystone in the theoretical framework known as the <u>Standard Model</u> of <u>particles</u> and their interactions. According to the Standard Model, the <u>Higgs boson</u> explains why some elementary particles have mass and others do not.

So far, the Higgs particle has eluded direct detection. Searches at the Large Electron Positron <u>collider</u> at the European laboratory <u>CERN</u> established that the Higgs boson must weigh more than  $114 \text{ GeV/c}^2$ . Calculations of quantum effects involving the Higgs boson require its mass to be less than  $185 \text{ GeV/c}^2$ .

So far, CDF and DZero each have analyzed about three inverse femtobarns of collision data--the scientific unit that scientists use to count the number of collisions. Each experiment expects to receive a total of about 10 inverse femtobarns by the end of 2010, thanks to the superb performance of the Tevatron. The collider continues to set numerous performance records, increasing the number of protonantiproton collisions it produces.

Provided by Fermi National Accelerator Laboratory



Citation: New experiments constrain Higgs mass (w/Videos) (2009, March 13) retrieved 4 May 2024 from <u>https://phys.org/news/2009-03-constrain-higgs-mass-wvideos.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.