

# Hubble provides new evidence for dark matter around small galaxies

March 12 2009

---



These four dwarf galaxies are part of a census of small galaxies in the tumultuous heart of the nearby Perseus galaxy cluster. The galaxies appear smooth and symmetrical, suggesting that they have not been tidally disrupted by the pull of gravity in the dense cluster environment. Larger galaxies around them, however, are being ripped apart by the gravitational tug of other galaxies. The images, taken by NASA/ESA's Hubble Space Telescope, are evidence that the undisturbed galaxies are enshrouded by a "cushion" of dark matter, which protects them from their rough-and-tumble neighborhood. Dark matter is an invisible form of matter that accounts for most of the Universe's mass. Astronomers have deduced the existence of dark matter by observing its gravitational influence on normal matter, consisting of stars, gas, and dust. Observations by Hubble's Advanced Camera for Surveys spotted 29 dwarf elliptical galaxies in the Perseus Cluster, located 250 million light-years away and one of the closest galaxy clusters to Earth. Of those galaxies, 17 are new discoveries. The images were taken in 2005. Credit: NASA, ESA and C.

Conselice (University of Nottingham, U.K.)

Peering into the tumultuous heart of the nearby Perseus galaxy cluster, Hubble discovered a large population of small galaxies that have remained intact while larger galaxies around them are being ripped apart by the gravitational tug of other galaxies.

The Hubble images provide further evidence that the undisturbed [galaxies](#) are enshrouded by a "cushion" of [dark matter](#) that protects them from their rough-and-tumble neighbourhood.

Dark matter is an invisible form of matter that accounts for most of the Universe's mass. Astronomers have deduced the existence of dark matter by observing its gravitational influence on normal matter, such as stars, gas and dust.

"We were surprised to find so many [dwarf galaxies](#) in the core of this cluster that were so smooth and round and had no evidence at all of any kind of disturbance," says astronomer Christopher Conselice of the [University of Nottingham](#), UK, and leader of the team that made the Hubble observations. "These dwarfs are very old galaxies that have been in the cluster for a long time. So if something was going to disrupt them, it would have happened by now. They must be very, very dark-matter-dominated galaxies."

The dwarf galaxies may have an even higher amount of dark matter than spiral galaxies. "With these results, we cannot say whether the dark matter content of the dwarfs is higher than in the Milky Way Galaxy," Conselice says. "Although, the fact that spiral galaxies are destroyed in clusters, while the dwarfs are not, suggests that this is indeed the case."

First proposed about 80 years ago by Swiss astronomer Fritz Zwicky, dark matter is thought to be the glue that holds galaxies together. Astronomers suggest that dark matter provides a vital scaffolding for the Universe, forming a framework for the [formation of galaxies](#) through gravitational attraction.

Observations by Hubble's Advanced Camera for Surveys spotted 29 dwarf [elliptical galaxies](#) in the Perseus Cluster, located 250 million light-years away and one of the closest galaxy clusters to Earth. Of these galaxies, 17 are new discoveries.

Because dark matter cannot be seen, astronomers detected its presence through indirect evidence. The most common method is by measuring the velocities of individual stars or groups of stars as they move randomly in the galaxy or as they rotate around the galaxy. The Perseus Cluster is too far away for telescopes to resolve individual stars and measure their motions. So Conselice and his team derived a new technique for uncovering dark matter in these dwarf galaxies by determining the minimum additional mass contribution from dark matter that the dwarfs must have to protect them from being disrupted by the strong, tidal pull of gravity from larger galaxies.

Studying these [small galaxies](#) in detail was possible only because of the sharpness of Hubble's Advanced Camera for Surveys. Conselice and his team first spied the galaxies with the WIYN Telescope at Kitt Peak National Observatory. These observations, Conselice says, only hinted that many of the galaxies were smooth and therefore dark matter dominated. "Those ground-based observations could not resolve the galaxies, so we needed Hubble imaging to nail it," he says.

The Hubble results appeared in the 1 March issue of *Monthly Notices of the Royal Astronomical Society*.

Source: ESA/Hubble Information Centre ([news](#) : [web](#))

Citation: Hubble provides new evidence for dark matter around small galaxies (2009, March 12)  
retrieved 25 April 2024 from

<https://phys.org/news/2009-03-hubble-evidence-dark-small-galaxies.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.