

# Swift Satellite records early phase of gamma ray burst

March 2 2009

---



Illustration of GRB. Credit:NASA

(PhysOrg.com) -- UK astronomers, using a telescope aboard the NASA Swift Satellite, have captured information from the early stages of a gamma ray burst - the most violent and luminous explosions occurring in the Universe since the Big Bang. The work was published on Friday 27th February in the *Monthly Notices of the Royal Astronomical Society*.

Swift is able to both locate and point at gamma ray bursts (GRBs) far quicker than any other telescope, so by using its Ultraviolet/Optical Telescope (UVOT) the astronomers were able to obtain an ultraviolet spectrum of a GRB just 251 seconds after its onset - the earliest ever captured. Further use of the instrument in this way will allow them to calculate the distance and brightness of GRBs within a few hundred

seconds of their initial outburst, and gather new information about the causes of bursts and the galaxies they originate from.

It is currently thought that some GRBs are caused by immense explosions following the collapse of the core of a rapidly rotating, high-mass star into a black hole, but there are still many mysteries surrounding them.

"The UVOT's wavelength range, coupled with the fact that Swift is a space observatory with a speedy response rate, unconstrained by time of day or weather, has allowed us to collect this early ultraviolet spectrum," said Martin Still from the Mullard Space Science Laboratory (MSSL) at UCL.

Paul Kuin, also from MSSL, who works on the calibration of the UVOT instrument explained: "By looking at these earlier moments of gamma ray bursts, we will not only be able to better calculate things such as the luminosity and distance of a burst, but to find out more about the galaxies that play host to them and the impact these explosions have on their environments. Once this new technique is applied to much brighter bursts, we'll have a wealth of new data."

Massimiliano De Pasquale, a GRB scientist of the UVOT team from MSSL, added, "The UVOT instrument is particularly suited to study bursts with an average to high redshift (1) - a part of the ultraviolet spectrum that is difficult for even the very big ground-based telescopes to study. Using UVOT with Swift, we can now find redshifts for bursts that were difficult to capture in the past and find out more about their distant host galaxies, about ten billion light years away."

Professor Keith Mason, Chief Executive of the Science and Technology Facilities Council, said, "This is an amazing first for the UVOT instrument and an exciting new development in the study of these most

violent and energetic explosions. Thanks to the hard work of our UK scientists at MSSL, and their partners, we can now gather far more information about gamma ray bursts and the early Universe."

Since its launch in 2004, the Swift satellite has provided the most comprehensive study so far of GRBs and their afterglows. Using the UVOT to obtain ultraviolet spectrums, the Swift team will be able to build on this study and even determine more about the host galaxies' chemistry.

Paul Kuin said, "The new spectrum has not only allowed us to determine the distance of the gamma ray burst's host galaxy but has revealed the density of its hydrogen clouds. Learning more about these far-away galaxies helps us to understand how they formed during the early universe. The gamma ray burst observed on this occasion originated in a galaxy 8 billion light years from Earth."

## **Note:**

(1) Redshift is the systematic displacement of individual lines in the spectrum of a celestial object toward the red or longer wavelength, end of the visible spectrum. All distant galaxies show a redshift proportional to their distance from the earth as a result of the general expansion of space. Known as the cosmological red shift, this results when the wavelength of light is stretched as it moves through the expanding universe.

## **Swift**

Launched in November 2004, Swift detects gamma-ray burst and X-ray flashes, and relays their coordinates to the scientific community within seconds. While Swift rapidly re-points to continue observations with its

high resolution telescopes of the decaying X-ray, UV and optical afterglows that accompany these events, rapid follow-up observations may also be undertaken by other more powerful satellites such as Chandra, the Hubble Space Telescope or XMM satellites and by the European Southern Observatory's Very Large Telescope (VLT) and other major observatories. These powerful co-ordinated observation programmes, instigated by Swift, are slowly unravelling the mystery of star explosions.

Source: Science and Technology Facilities Council

Citation: Swift Satellite records early phase of gamma ray burst (2009, March 2) retrieved 15 August 2024 from <https://phys.org/news/2009-03-swift-satellite-early-phase-gamma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.